

A Revised List of Bryophytes Reported from Seram (Ceram) and Ambon Islands, the Moluccas, Based on Collections Made on the 1984–1986 Japan-Indonesia Collaborative Expeditions

HIROYUKI AKIYAMA

*Museum of Nature and Human Activities, Hyogo, Yayoigaoka-6, Sanda, Hyogo 669-1546, Japan.
akiyama@hitohaku.jp*

A total of 610 taxa of bryophytes (48 families, 154 genera, 364 species, and 3 varieties in mosses, 21 families, 59 genera, 226 species, 7 varieties and 1 forma in liverworts, and 1 family, 4 genera, and 9 species in hornworts) are reported from Ambon and Seram Islands based on collections made on the 1984–1986 Japan-Indonesian collaborative expeditions.

Keywords: Ambon, bryophytes, Ceram, flora, hepatics, hornworts, Moluccas, mosses, Seram

Seram (or Ceram) and the Ambon Islands are situated in the central Moluccas in eastern Maleisia between two large islands, Sulawesi and New Guinea. Because of their location, it is expected that the bryophyte flora of these islands would show an intermediate feature between Malesian and Melanesian elements.

I have had a chance to conduct bryological surveys on both islands, mainly on Seram, as a member of the Japan-Indonesian collaborative expeditions from 1984 to 1986 (Fig. 1). Seram is composed mostly of limestone, especially in the central part, where the Murkele Ridge, which has several peaks exceeding 2500 m, with the highest being Gunung (Mt.) Binaiya at 3027 m, is located. In eastern and western Seram are low mountains not exceeding 1500 m (mostly lower than 500 m). The central part of the island now comprises Manusela National Park.

Before our research activities, the bryophyte flora of Seram and Ambon islands was known only from a few reports by Bartram (1939), Dozy & Molkenboer (1844, 1845–1854, 1854–1861, 1861–1870), Fleischer (1900–1922), Herzog (1916,

1919, 1921, 1926), Schiffner (1893, 1910), and Stephani (1898–1924). Additional papers on the bryophyte flora of Ambon and Seram, which are totally or partially based on our recent collections, as follows: Akiyama (1986, 1988, 1989, 1990, 1992, 1993, 1996, 1997a, b, 1999, 2006, 2007, 2009), Akiyama & Reese (1993), Akiyama & Suleiman (2001), Akiyama *et al.* (1991), Câmara (2009), Doei (1987a, b), Hattori (1986), Hasegawa (1986), Inoue (1986), Iwatsuki & Suzuki (1988), Kitagawa (1986), Klazenga (1999), Kruijer (2002), Mizutani (1986), Reese *et al.* (1991), Touw (1993, 2001a, b), Vitt (1995), Yamaguchi (1993), Watanabe & Akiyama (1991) and van Zanten (2006).

As a result, 48 families, 154 genera, 364 species, and 3 varieties of mosses, 21 families, 59 genera, 226 species, 7 varieties, and 1 forma of liverworts, and 1 family, 4 genera, and 9 species of hornworts are recognized among our collections from these two islands. Eight species of mosses, 11 species and one variety of hepatics, and one species of hornworts were described as new based on the collections (Table 1). Forty-




FIG. 1. Ambon and Seram Islands. Central part of the Seram Island is divided into four parts, North, Central, South, and the Manusela National Park (shaded).

three species in 33 genera of mosses and 15 species in eleven genera of hepatics, which were reported from the two islands (mostly from Ambon, where nearly all natural vegetation has been destroyed on Ambon because of intensive cultivation) in the older literature were not found during our expeditions.

TABLE 1. Taxa from Seram and Ambon islands described as new based on collections made on the 1984–1986 Japan-Indonesia collaborative expeditions.

Mosses

- Barbula seramense* H. Akiyama
- Calomnion ceramensis* Vitt
- Calypstrohaeta isophylla* H. Akiyama
- Holomitrium seramensis* H. Akiyama
- Leskeodon seramensis* H. Akiyama
- Powellia macromitrioides* H. Akiyama
- Syrrhopodon pulcher* Reese
- Taxithelium damanhurianum* P.S. Câmara

Liverworts

- Frullania akiyamae* S. Hatt.
- F. cordistipula* (Reinw. et al.) Dumort.
var. *dentistipula* S. Hatt.
- F. gracilis* (Reinw. et al.) Dumort. var. *vittata* S. Hatt.
- F. hasskarliana* Lindenb. var. *gracilis* S. Hatt.
- F. hasskarliana* Lindenb. var. *parvidentata* S. Hatt.
- F. ptiophylla* S. Hatt. var. *minor* S. Hatt.
- F. serrata* Gottsche var. *ceramensis* S. Hatt.
- F. suocellata* S. Hatt.
- Hattorirolejeunea akiyamae* Mizut.
- Lepidozia integrifolia* Doei
- Plagiochila akiyamae* Inoue
- P. ceramica* Inoue

Hornwort

- Dendroceros seramensis* J. Haseg.

Bryophytes of Seram and Ambon

In the list below, families, genera, and species are arranged alphabetically. Taxa newly described based on our collections are preceded by an asterisk (*).

Abbreviations of localities are as follows: **A** = Ambon, **WS** = western Seram, **CS** = central Seram (600–800 m, central inland areas excluding Manusela National Park), **SS** = southern Seram (below 500 m, southern inland areas excluding Manusela National Park), **NS** = northern Seram (0–600 m), **ES** = eastern Seram, **MNP** = Manusela National Park, central Seram (600–3000 m). For details on collecting sites in the islands, see figure 1 in this report and in Akiyama (1989, figures 2–4). Note that to determine the bryophyte flora of Manusela National Park, we tentatively divided central Seram into four subregions: CS, MNP, NS, and SS.

MUSCI

Amblystegiaceae

Platyhypnidium M. Fleisch.

- P. muelleri** (A. Jaeger) M. Fleisch. —Akiyama (1986: 62); Akiyama (1993: 229). MNP, SS.

Bartramiaceae

Bartramia Hedw.

- B. hallerana** Hedw. —Akiyama (2009). MNP.

Breutelia (Bruch & Schimp.) Schimp.

- B. arundinifolia** (Duby) M. Fleisch. —Akiyama (1996: 50); Akiyama (2009). MNP.

Leiomela (Mitt.) Broth.

- L. javanica** (Renauld & Cardot) Broth. —Akiyama (2009). MNP.

Philonotis Brid.

- P. calomicra** Broth. —Akiyama (2009). A, SS, WS.
P. hastata (Duby) Wijk & Margad. —Akiyama (2009). CS, SS, WS.
P. roylei (Hook.f.) Mitt. —Akiyama (2009). WS.
P. secunda (Dozy & Molk.) Bosch & Sande Lac. —Akiyama (2009). CS, MNP.
P. turneriana (Schwägr.) Mitt. —Akiyama (2009). MNP, SS, WS.

Brachytheciaceae

Palamocladium Müll. Hal.

- P. nilgheriense* (Mont.) Müll. Hal. = *P. leskeoides*, fide Hofmann (1997).
P. leskeoides (Hook.) E. Britton —Akiyama (1993: 229, as *P. nilgheriense*); Akiyama (2009). MNP.

Bryaceae

Anomobryum Schimp.

- A. julaceum** (Schrader ex P. Gaertn. *et al.*) Schimp. —Akiyama (2009). MNP.

Brachymenium Schwägr.

- B. exile** (Dozy & Molk.) Bosch & Sande Lac. —Akiyama (2009). SS.
B. gracile E.B. Bartram —Akiyama (2009). MNP.
B. indicum (Dozy & Molk.) Bosch & Sande Lac. —Akiyama (2009). SS.
B. nepalense Hook. —Akiyama (2009). MNP.
B. cf. nepalense Hook. —Akiyama (2009). MNP.

Bryum Hedw.

- B. apiculatum** Schwägr. —Akiyama (2009). MNP, SS, WS.
B. argenteum Hedw. —Akiyama (2009). MNP.
B. billardieri Schwägr. —Akiyama (2009). MNP.
B. cf. bicolor —Akiyama (2009). MNP.
B. clavatum (Schimp.) Müll. Hal. —Akiyama (2009). SS.
B. leucophyllum Dozy & Molk. —Akiyama (2009). MNP.
B. perdecurrens E.B. Bartram —Akiyama (2009). MNP.
B. tuberosum Mohamed & Damanhuri —Akiyama (2009). MNP.

Leptodontium (Müll. Hal.) Hampe ex Lindb.

- L. flexifolium** (With.) Hampe —Akiyama (2009). MNP.

Orthodontium Schwägr.

- O. inflatum** Dozy & Molk. —Akiyama

(2009). MNP, WS.

Rhodobryum (Schimp.) Limpr.

R. aubertii (Schwägr.) Thér. —Akiyama (2009). CS, NS, SS, WS.

R. giganteum (Schwägr.) Paris —Akiyama (2009). MNP.

Buxbaumiaceae

Buxbaumia Hedw.

B. javanica Müll. Hal. —Akiyama (1986: 39); Akiyama (1989: 151). MNP.

Calomniaceae

Calomnion Hook.f. & Wils.

C. ceramense Vitt —Akiyama (1988: 12, as *C. complanatum*); Akiyama (1990: 401, as *C. cf. complanatum*); Vitt (1995: 339); Akiyama (2009). MNP.

Type: Manusela National Park, Tehoru, Nihata to summit of Gunung Hoale Besar, 1560 m alt., *Akiyama C-16703* (holotype in KYO; isotypes in FH, NY).

C. complanatum (Hook.f. & Wils.) Lindb. = misidentification of *C. ceramense*, fide Vitt (1995: 339).

C. cf. complanatum (Hook.f. & Wils.) Lindb. = misidentification of *C. ceramense*, fide Vitt (1995: 339) and Akiyama (2009).

Calymperaceae

Arthrocormus Dozy & Molk.

A. shimperi (Dozy & Molk.) Dozy & Molk. —Yamaguchi (1993b: 301). CS, ES, NS, SS, WS.

Calymperes Sw. ex F. Weber

C. aeruginosum Hampe ex Sande Lac. —Akiyama (1986: 44); Akiyama & Reese (1993: 200). CS, MNP, NS, SS, WS.

C. boulayi Besch. —Akiyama & Reese (1993: 201). ES, SS, WS.

C. couguiense Besch. —Akiyama & Reese (1993: 201). SS.

C. erosum Müll. Hal. —Akiyama & Reese (1993: 202). A, ES, WS.

C. graeffeanum Müll. Hal. —Akiyama & Reese (1993: 202). ES, NS, SS, WS.

C. lonchophyllum Schwägr. —Akiyama

(1986: 44, as *C. aff. serratum*); Akiyama & Reese (1993: 202). A, ES, NS, SS, WS.

C. longifolium Mitt. = misidentification of *Syrrhpodon loreus* (Calymperaceae), fide Akiyama & Reese (1993: 209).

C. palisotii Schwägr. subsp. **moluccense** (Schwägr.) M. Menzel —Akiyama & Reese (1993: 203). NS, SS, WS.

C. porrectum Mitt. —Akiyama & Reese (1993: 203). A, NS, SS, WS.

C. serratum A. Braun ex Müll. Hal. —Akiyama (1986: 44); Akiyama & Reese (1993: 204). NS, MNP, SS, WS.

C. aff. serratum = misidentification of *C. lonchophyllum*, fide Akiyama & Reese (1993: 202).

C. strictifolium (Mitt.) G. Roth —Akiyama (1986: 44); Akiyama & Reese (1993: 205). A, NS, CS, SS.

C. subintegrum Broth. —Akiyama & Reese (1993: 205). A, ES.

C. taitense (Sull.) Mitt. —Akiyama & Reese (1993: 205). A, CS, ES, MNP, NS, SS, WS.

C. trenerum Müll. Hal. —Akiyama & Reese (1993: 206). SS, WS.

Exodictyon Cardot

E. incrassatum (Mitt.) Cardot ex Paris —Yamaguchi (1993b: 301). SS.

Exostratum L.T. Ellis

E. asperum (Mitt.) L.T. Ellis —Yamaguchi (1993b: 301). NS.

E. blumei (Nees ex Hampe) L.T. Ellis —Yamaguchi (1993b: 302). MNP, NS, SS.

E. sullivanii (Dozy & Molk.) L.T. Ellis —Yamaguchi (1993b: 302). MNP.

Leucophanes Brid.

L. albescens Müll. Hal. = *L. glaucum*, fide Salazar Allen (1993).

L. angustifolium Renauld & Cardot —Akiyama (1986: 43, as *L. massartii*); Yamaguchi (1993b: 302). CS, ES, MNP, NS, SS, WS.

L. candidum (Schwägr.) Lindb. —Akiyama (1986: 43); Yamaguchi (1993b: 302). CS, MNP, NS, SS, WS.

L. glaucum (Schwägr.) Mitt. —Akiyama (1986: 43, as *L. albescens*); Yamaguchi

(1993b: 302). A, ES, NS, SS, WS.

L. massartii Renauld & Cardot = *L. angustifolium*, fide Salazar Allen (1993).

L. octoblepharioides Brid. —Akiyama (1986: 43); Yamaguchi (1993b: 302). A, ES, MNP, NS, SS, WS.

Mitthyridium Robins.

M. constrictum (Sull.) Robinson —Akiyama (1986: 45); Akiyama & Reese (1993: 213). A, MNP, SS, WS.

M. crassum (Hook. & Grev.) H. Robins. —Akiyama (1986: 45, as *M. cf. crassum*); Akiyama & Reese (1993: 213). WS.

M. cf. crassum (Hook. & Grev.) H. Robins. = misidentification of *M. crassum*, fide Akiyama & Reese (1993: 213).

M. fasciculatum (Hook. & Grev.) H. Robins. —Akiyama (1986: 45, as *M. cf. fasciculatum*); Akiyama & Reese (1993: 213). A, ES, MNP, NS, SS, WS.

M. cf. fasciculatum (Hook. & Grev.) H. Robins. = misidentification of *M. fasciculatum*, fide Akiyama & Reese (1993: 213).

M. leucoloma (Müll. Hal.) H. Robins. —Akiyama & Reese (1993: 214). ES.

M. louisiadum (Broth.) H.K. Nowak = *M. repens*, fide Reese (1986).

M. papuanum (Broth.) H. Robins. —Akiyama & Reese (1993: 214). NS.

M. perundulatum (Broth.) H. Robins. —Akiyama & Reese (1993: 215). NS, WS.

M. repens (Harv.) H. Robins. —Akiyama (1986: 45, as *M. louisiadum*); Akiyama & Reese (1993: 215). ES, NS, SS, WS.

M. subluteum (Müll. Hal.) Nowak. —Akiyama (1986: 45); Akiyama & Reese (1993: 215). CS, MNP, NS, SS, WS.

M. undulatum (Dozy & Molk.) H. Robins. —Akiyama & Reese (1993: 216). CS, MNP, NS, SS, WS.

M. wallisii (Müll. Hal.) H. Robins. —Akiyama & Reese (1993: 216). SS.

Octoblepharum Hedw.

O. albidum Hedw. —Akiyama (1986: 43); Yamaguchi (1993b: 303). CS, ES, MNP, NS, WS.

Syrhophodon Schwägr.

S. albovaginatus Schwägr. —Akiyama & Reese (1993: 207). A, ES, NS, SS, WS.

S. aristifolius Mitt. —Akiyama & Reese (1993: 207). SS, WS.

S. banksii Müll. Hal. = *S. involutus*, fide Mohamed & Reese (1985).

S. ciliatus (Hook.) Schwägr. —Akiyama (1986: 45, as *S. cf. ciliatus*); Akiyama & Reese (1993: 208). NS, SS, WS.

S. cf. ciliatus (Hook.) Schwägr. = misidentified as *S. ciliatus*, fide Akiyama & Reese (1993: 208).

S. croceus Mitt. —Akiyama & Reese (1993: 208). A, NS, SS, WS.

S. gardneri (Hook.) Schwägr. —Akiyama & Reese (1993: 208). CS, MNP.

S. hispidissimus Dixon —Akiyama & Reese (1993: 209). NS.

S. involutus Schwägr. —Akiyama (1986: 47, as *S. bankii*); Akiyama & Reese (1993: 209). SS, WS.

S. loreus (Sande Lac.) Reese —Akiyama (1986: 44, as *Calymperes longifolium*); Akiyama & Reese (1993: 209). NS, SS, WS.

S. muelleri (Dozy & Molk.) Sande Lac. —Akiyama & Reese (1993: 210). A, CS, MNP, NS, SS, WS.

***S. pulcher** Reese in Reese *et al.* —Reese *et al.* (1991: 419); Akiyama & Reese (1993: 210). MNP, WS.

Type: West Seram, Gunung Tiang Bendera near Tanahgoyang, 900 m alt., Akiyama C-15577 (holotype in KYO, isotypes in BO, L, LAF, NY).

S. spiculosus Hook. & Grev. —Akiyama & Reese (1993: 210). ES, MNP, WS.

S. trachyphyllus Mont. —Akiyama & Reese (1993: 211). MNP.

S. tristichus Nees ex Schwägr. —Akiyama & Reese (1993: 211). CS, MNP, NS, WS.

Cryphaeaceae

Forsstoremia Lindb.

Forsstroemia sp. = misidentification of *Neolindbergia vitiensis* (Pterobryaceae), fide Akiyama *et al.* (1991: 82).

Daltoniaceae

Archboldiella E.B. Bartram

- A. nymaniana** (M. Fleisch.) H. Akiyama — Akiyama (1986: 56, as *A. pilifera*); Akiyama (1986: 58, as *Distichophyllidium* cf. *nymanianum*); Akiyama (1990: 402). NS, SS.
A. pilifera E.B. Bartram = *A. nymanianum*, fide Akiyama (1990: 402).

Calypstrochaeta Desv.

- ***C. isophylla** H. Akiyama — Akiyama (1988: 58, as *Eriopus* sp.); Akiyama (1990: 403). CS, MNP, NS.

Type: Central Seram, Seram Utara, near Elemata-Makualaina, 400–650 m alt., Akiyama C-9396 (holotype in KYO).

- C. remotifolia** (Müll. Hal.) Z. Iwats. *et al.* — Akiyama (1986: 58, also as *Eriopus parviretis*); Akiyama (1988: 58, as *Eriopus remotifolia* and *E. parviretis*); Akiyama (1990: 403). MNP, SS.

Daltonia Hook. & Tayl.

- D. angustifolia** Dozy & Molk. — Akiyama (1988: 31); Akiyama (1990: 405). MNP.
D. contorta Müll. Hal. — Akiyama (1986: 58, as *D. cf. contorta*); Akiyama (1988: 31); Akiyama (1990: 404). MNP.
D. cf. contorta Müll. Hal. = misidentification of *D. contorta*, fide Akiyama (1988: 31).

Distichophyllidium M. Fleisch.

- D. jungermanniaceum** Müll. Hal. ex M. Fleisch. — Akiyama (1986: 58, both as *D. cf. jungermanniaceum* and *Distichophyllum* cf. *jungermannioides*); Akiyama (1988: 38, as *D. jungermannioides*); Akiyama (1990: 409, as *D. jungermannioides*). MNP, SS.
D. jungermannioides (Müll. Hal.) M. Fleisch. = misspelling of *D. jungermanniaceum*.
D. cf. jungermanniaceum Müll. Hal. ex M. Fleisch. = misidentification of *D. jungermanniaceum*, fide Akiyama (1990: 409).
D. cf. nymanianum M. Fleisch. = misidentification of *Archboldiella nymaniana* (Daltoniaceae), fide Akiyama (1990: 402).

Distichophyllum Dozy & Molk.

- D. cuspidatum** (Dozy & Molk.) Dozy & Molk. — Akiyama (1988: 34); Akiyama

(1990: 406). MNP, SS, WS.

D. cf. cuspidatum (Dozy & Molk.) Dozy & Molk. — Akiyama (1986: 59) = *Distichophyllum* sp.

D. denticulatum Dixon — Akiyama (1988: 35, as *D. aff. santosii*); Akiyama 1990: 408. MNP, SS.

D. jungermannioides (Müll. Hal.) Bosch & Sande Lac. — Akiyama (1990: 409). WS.

D. cf. jungermannioides (Müll. Hal.) Bosch & Sande Lac. = *Distichophyllum jungermanniaceum*, fide Akiyama (1990: 409).

D. laxirete Nog. & Z. Iwats. — Akiyama (1988: 35); Akiyama (1990: 407). MNP, SS.

D. aff. laxirete Nog. & Z. Iwats. = misidentification of *D. subnigricaula*.

D. mittenii Bosch & Sande Lac. — Akiyama (1986: 59, both as *D. cf. osterwaldii* and *D. aff. osterwaldii*); Akiyama (1988: 34); Akiyama (1990: 407). CS, MNP, SS.

D. cf. osterwaldii M. Fleisch. = misidentification of *D. mittenii*, fide Akiyama (1990: 407).

D. aff. osterwaldii M. Fleisch. = misidentification of *D. mittenii*, fide Akiyama (1990: 407).

D. rhizophorum (M. Fleisch.) Crosby — Akiyama (1988: 33); Akiyama (1990: 406). MNP.

D. rotundatum Nog. & Z. Iwats. — Akiyama (1990: 405). MNP, SS, WS.

D. aff. santosii E.B. Bartram = misidentification of *D. denticulatum*, fide Akiyama (1988: 408).

D. subnigricaula Broth. — Akiyama (1990: 404, as *D. aff. laxirete*). MNP, SS.

D. cf. subnigricaula Broth. = *D. subnigricaula*.

Ephemeropsis K.I. Goebel

E. tjibodensis K.I. Goebel — Akiyama (1986: 60); Akiyama (1988: 38); Akiyama (1990: 411). SS, WS.

Eriopus (Brid.) Brid.

E. parviretis M. Fleisch. = misidentification of *Calypstrochaeta remotifolia* (Daltoniaceae), fide Akiyama (1990: 403).

E. remotifolia Müll. Hal. = *Calypstrochaeta*

remotifolia.

Eriopus sp. = *Calyptrochaeta isophylla*.

Leskeodon Broth.

L. acuminatus (Bosch & Sande Lac.) M. Fleisch. —Akiyama (1986: 60, as *L. cf. acuminatus*); Akiyama (1988: 40); Akiyama (1990: 411). MNP, SS.

L. cf. acuminatus (Bosch & Sande Lac.) M. Fleisch. = misidentification of *L. acuminatus*, fide Akiyama (1990: 411).

L. aff. acuminatus —Akiyama (1986: 60).

Note. Akiyama (1990) did not include the species in his report on the genus.

***L. seramensis** H. Akiyama —Akiyama (1988: 40, as *Leskeodon* sp. 2); Akiyama (1990: 410). SS.

Type: Central Seram, Tehoru, Saunule-Wae-Waya, 110–190 m alt., Akiyama C-14646 (holotype in KYO). SS.

Dicranaceae

Atractylolcarpus Mitt.

A. comosus Dixon —Akiyama (1990: 386). MPN.

Braunfelsia Paris

B. dicranoides (Dozy & Molk.) Broth. —Akiyama (1986: 40, as *B. aff. dicranodites*); Akiyama (1990: 393).

B. aff. dicranoides (Dozy & Molk.) Broth. = misidentification of *B. dicranodites*, fide Akiyama (1990: 393).

B. plicata (Sande Lac.) Broth. —Akiyama (1990: 394).

B. edentula (Mitt.) Wijk & Margad. —Akiyama (1990: 394).

Brotherobryum M. Fleisch.

B. sleumeri B. Allen —Akiyama (2009). MNP.

Bryohumbertia P. de la Varde & Thér.

B. walkeri (Mitt.) J.-P. Frahm —Akiyama (1990: 386). MPN.

Campylopodium (Müll. Hal.) Besch.

C. medium (Duby) Giese & J.P. Frahm —Akiyama (1990: 386). WS.

Campylopus Brid.

C. aureus Bosch & Sande Lac. —Akiyama (1990: 388). MNP.

C. clemensiae E.B. Bartram —Akiyama (1990: 388). MNP.

C. exasperatus (Nees & Blume) Brid. var. **lorentzii** (M. Fleisch.) J.P. Frahm —Akiyama (1990: 387). MPN.

C. exasperatus (Nees & Blume) = misidentification of *C. exasperatus* var. *lorentzii*, fide Akiyama (1990: 387).

C. involutus (Müll. Hal.) A. Jaeger —Akiyama (1990: 387). WS.

Cryptodicranum E.B. Bartram

C. armitii (Müll. Hal.) E.B. Bartram —Akiyama (1986: 41); Akiyama (1990: 394). MNP.

Dicranella (Müll. Hal.) Schimp.

D. coarctata (Müll. Hal.) Bosch & Sande Lac. —Akiyama (1990: 390). A, MNP.

D. setifera (Mitt.) A. Jaeger —Akiyama (1986: 40); Akiyama (1990: 389). MNP, WS.

Dicranodontium Bruch & Schimp.

D. fleischerianum Schultz-Motel —Akiyama (1986: 40); Akiyama (1990: 387). MNP.

Dicranoloma (Renauld) Renauld

D. assimile (Hampe) Renauld —Akiyama (1990: 398); Klazenga (1999: 54). MNP.

D. billardierei (Brid. ex Anon.) Paris —Klazenga (1999: 63). MNP.

D. blumii (Nees) Paris —Akiyama (1986: 40); Akiyama (1990: 395); Klazenga (1999: 69). MNP, WS.

D. braunii (Müll. Hal.) Paris —Akiyama (1990: 399); Klazenga (1999: 76). MNP, NS, WS.

D. brevisetum (Dozy & Molk.) Paris —Akiyama (1986: 40); Akiyama (1990: 400, as *D. leucophyllum*); Klazenga (1999: 82). MNP.

D. cutlackii (Norris & T.J. Kop.) Klazenga —Akiyama (1990: 386, as *D. dymannianum*); Klazenga (1999: 86); Akiyama (2009). MNP.

D. dymannianum E.B. Bartram = misidentification of *D. cutlackii*, fide Klazenga (1999: 87).

D. leucophyllum (Hampe ex Sande Lac.) Paris = *D. brevisetum*, fide Tan & Koponen

(1983) and Klanzega (1999: 79); also see the note given by Klazenga (1999: 83).

Holomitrium Brid.

H. cylindraceum (P. Beuav.) Wijk & Margad. —Akiyama (1993: 250). SS.

***H. seramense** H. Akiyama —Akiyama (1990: 393). NNP.

Type: Central Seram, Kecamatan Seram Utara, en route from Wae Huhu to Gunung O'Wae Puku, 2080 m alt., *Akiyama* C-8863 (holotype in KYO).

Leucoloma Brid.

L. molle (Müll. Hal.) Mitt. —Akiyama (1986: 41); Akiyama (1990: 388). MNP, SS, WS.

Microdus Schimp. ex Besch.

M. brasiliensis (Duby) Thériot —Akiyama (1990: 391). WS.

M. cf. insularis (Willams) Broth. = misidentification of *M. miquelianus*, fide Akiyama (1990: 391).

M. aff. leibergii (Willams) Broth. = misidentification of *M. miquelianus*, fide Akiyama (1990: 391).

M. miquelianus (Mont.) Besch. in Paris —Akiyama (1990: 391). MNP, NS, SS, WS.

M. cf. pomiformis (Griff.) Besch. = misidentification of *M. miquelianus*, fide Akiyama (1990: 391).

Trematodon Michx.

T. longicollis Michx. —Akiyama (1986: 42, as *T. aff. longicollis*); Akiyama (1990: 389). WS.

T. aff. longicollis Michx. = misidentification of *T. longicollis*, fide Akiyama (1990: 389).

Wilsoniella Müll. Hal.

W. decipiens (Mitt.) Alston var. **acutifolia** (Dixon) Wijk. & Margad. —Akiyama (1986: 42, as *W. pellucida*); Akiyama (1990: 389, as *W. pellucida*). WS.

W. pellucida (Wils.) Müll. Hal. = *Wilsoniella decipiens* var. *acutifolia*, fide Chien (1994).

Diphysciaceae

Diphyscium Mohr.

D. longifolium Griff. —Akiyama (1989: 152, as *D. loriae*). CS, MNP, SS, WS.

D. mucronifolium Mitt. —Akiyama (1986:

39, as *D. involutum*); Akiyama (1989: 152, as *D. involutum*). CS, SS, WS.

D. loriae Müll. Hal. = *D. longifolium*, fide Magonbo (2003).

D. involutum Mitt. = *D. mucronifolium*, fide Magonbo (2003).

Ditrichaceae

Ditrichum Hampe

D. difficile (Duby) M. Fleisch. —Akiyama (1986: 39); Akiyama (2009). MNP.

D. aff. colijnii Dixon —Akiyama (2009). CS, SS.

Garckeia Müll. Hal.

G. comosa (Dozy & Molk.) Wijk & Margad. = *G. flexuosa*, fide Norris & Koponen (1988).

Garckeia flexuosa (Griff.) Margad. & Nork. —Akiyama (1986: 40, as *G. comosa*); Akiyama (2009). WS.

Drepanophyllaceae

Mniomalia Müll. Hal.

M. semilimbata (Mitt.) Müll. Hal. —Akiyama (1986: 48); Akiyama (1989: 157). ES, SS.

Entodontaceae

Entodon Müll. Hal.

Entodon sp. —Akiyama (1986: 62).

Erpodiaceae

Solmsiella Müll. Hal.

S. biseriata (Austin) Steere —Akiyama (2009). WS.

Fissidentaceae

Fissidens Hedw.

F. bogoriensis M. Fleisch. —Iwatsuki & Suzuki (1988: 216). SS.

F. bryoides Hedw. var. **schimidii** (Müll. Hal.) R.S. Chopra & S.S. Kumar —Iwatsuki & Suzuki (1988: 216). MNP.

F. ceylonensis Dozy & Molk. —Iwatsuki & Suzuki (1988: 216). ES, MNP, WS.

F. crassinervis Sande Lac. —Iwatsuki & Suzuki (1988: 216). NS, WS.

F. crenulatus Mitt. —Iwatsuki & Suzuki

- (1988: 217). ES, SS, WS.
- F. crsipulus** Brid. —Iwatsuki & Suzuki (1988: 219, as *F. zippelinaus*). ES, SS, WS.
- F. flabellulus** Thwait. & Mitt. —Iwatsuki & Suzuki (1988: 217). MNP, WS.
- F. geminiflorus** Dozy & Molk. —Iwatsuki & Suzuki (1988: 217). MNP.
- F. hollianus** Dozy & Molk. —Iwatsuki & Suzuki (1988: 217). ES, MNP, SS, WS.
- F. hyallinus** Hook.f. & Wils. —Iwatsuki & Suzuki (1988: 217). MNP.
- F. laxus** Sull & Lesq. —Iwatsuki & Suzuki (1988: 217). ES, WS.
- F. microcladus** Thwait. & Mitt. —Iwatsuki & Suzuki (1988: 217). ES, MNP, WS.
- F. nobilis** Griff. —Iwatsuki & Suzuki (1988: 217). CS, ES, MNP, NS, SS, WS.
- F. oblongifolius** Hook.f. & Wils. —Iwatsuki & Suzuki (1988: 217). MNP, NS, MNP.
- F. pallidus** Hook.f. & Wils. —Iwatsuki & Suzuki (1988: 218). CS, MNP, SS, WS.
- F. punctulatus** Sande Lac. —Iwatsuki & Suzuki (1988: 218). CS, NS, MNP, SS.
- F. robinsonii** Broth. —Iwatsuki & Suzuki (1988: 218). ES, NS, SS, WS.
- F. serratus** Müll. Hal. —Iwatsuki & Suzuki (1988: 219). MNP.
- F. splachnobryoides** Broth. in Schum. —Iwatsuki & Suzuki (1988: 219). WS.
- F. subbryoides** Gangulee —Iwatsuki & Suzuki (1988: 219). WS.
- F. taxifolius** Hedw. —Iwatsuki & Suzuki (1988: 219). MNP.
- F. zollingeri** Mont. —Iwatsuki & Suzuki (1988: 219). MNP, NS, WS.
- F. zippelinaus* Dozy & Molk. = *Fissidens crsipulus*, fide Bruggeman-Nannenga (1997).

Garovagliaceae

Garovaglia Endl.

- G. angustifolia** Mitt. var. **bogoriense** (M. Fleisch.) During —Akiyama (1990: 140). MNP, NS, SS, WS.
- G. aristata** Bosch & Sande Lac. —Akiyama (1990: 140). NS, SS.
- G. baeuerlenii** (Geh.) Paris —Akiyama

- (1986: 54); Akiyama (1990: 140). NS, MNP.
- G. compressa** Mitt. var. **compressa** —Akiyama (1990: 140). SS, WS.
- G. aff. compressa* Mitt. = misidentification of *G. elegans*, fide Akiyama (1990: 141).
- G. cf. crispa** (Broth. & Geh.) Broth. —Akiyama (1990: 140). MNP.
- G. elegans** (Dozy & Molk.) Bosch & Sande Lac. —Akiyama (1986: 54, as *G. aff. compressa*); Akiyama (1986: 54); Akiyama (1990: 140). MNP, NS, SS, WS.
- G. elegans* (Dozy & Molk.) Bosch & Sande Lac. f. *elegans* = misidentification of *G. zantenii*, fide Akiyama (1990: 141).
- G. plicata** (Brid.) Bosch & Sande Lac. —Akiyama (1990: 140). MNP, WS.
- G. zantenii** During —Akiyama (1986: 54, as *G. elegans* f. *elegans*); Akiyama (1990: 140). MNP.

Hookeriaceae

Acrophyllum Vitt & Crosby

- A. javense** (Dixon ex Froelich) Z. Iwats. *et al.* —Akiyama (1988: 56); Akiyama (1997: 12). MNP.

Bryobrothera Thér.

- B. crenulata** (Broth. & Paris) Thér. —Akiyama (1997: 12). WS.

Callicostella (Müll. Hal.) Mitt.

- C. armata** Herzog —Akiyama (1988: 56, as *C. aff. papillata*); Akiyama (1997: 14). CS, MNP, SS, WS.
- C. papillata* (Mont.) Mitt. = misidentification of *C. probaktiana*, fide Akiyama (1997: 14).
- C. aff. papillata* (Mont.) Mitt. = misidentification of *C. armata*, fide Akiyama (1997: 14).
- C. probaktiana** (Müll. Hal.) Bosch & Sande Lac. —Akiyama (1986: 56, as *C. papillata*); Akiyama (1997: 14). MNP, NS, SS, WS.

Cyclodictyon Mitt.

- C. blumeinum** (Müll. Hal.) O. Kuntze —Akiyama (1986: 57); Akiyama (1988: 57); Akiyama (1997: 14). MNP, SS.

Hylocomiaceae

Macrothamnium M. Fleisch.

- M. javense** M. Fleisch. —Akiyama (1988: 52); Akiyama (2009). MNP.

Hypnaceae

Note. Although total nine species of *Ectropothecium* have been recognized among our collections (Akiyama 1988), they are still left in unidentified condition.

Ctenidiadelphus M. Fleisch.

- C. spinulus** M. Fleisch. —Akiyama (1988: 48); Akiyama (2009). NS.

Ctenidium (Schimp.) Mitt.

- C. malacobolum** (Müll. Hal.) Broth. —Akiyama (1988: 48); Akiyama (2009). MNP.
C. polychaetum (Bosch & Sande Lac.) Broth. —Akiyama (1988: 48); Akiyama (2009). MNP.

Ectropotheciella M. Fleisch.

- E. distichophylla** (Hampe ex Dozy & Molk.) M. Fleisch. —Akiyama (1988: 48); Akiyama (2009). CS, NS.

Hypnum Hedw.

- H. cf. plumaeforme** Wils. —Akiyama (1988: 51). MNP.

Isopterygium Mitt.

- I. cf. pohliaecarpum** (Sull. & Lesq.) A. Jaeger —Akiyama (1988: 51); Akiyama (2009). WS.

Vesicularia (Müll. Hal.) Müll. Hal.

- V. dubyana** (Müll. Hal.) Broth. —Akiyama (1988: 52, as *V. cf. dubyana*); Akiyama (2009). CS, NS.
V. cf. dubyana (Müll. Hal.) Broth. —misidentification of *V. dubyana*. fide Akiyama (2009).
V. motagnei (Bel.) M. Fleisch. —Akiyama (1988: 52, as *V. cf. monatnei*); Akiyama (2009). NS, WS.
V. cf. motagnei (Bel.) M. Fleisch. —misidentification of *V. montagnei*. fide Akiyama (2009).
V. reticulata (Dozy & Molk.) Broth. —Akiyama (1988: 52, as *V. cf. reticulata*); Akiyama (2009). NS, SS.
V. cf. reticulata (Dozy & Molk.) Broth. —

misidentification of *V. reticulata*. fide Akiyama (2009).

Stereodontopsis R.S. Williams

- S. excavata** (Broth.) Ando —Akiyama (2009). NS.

Hypnodendraceae

Hypnodendron Mitt.

- H. auricomum** Broth. & Geh. subsp. **auricomum** —Akiyama (1986: 49); Akiyama (1988: 19); Akiyama (1989a: 201); Akiyama (1989b: 159). MNP.
H. dendroides (Brid.) Touw —Akiyama (1986: 49); Akiyama (1988: 19); Akiyama (1989b: 158). MNP, WS.
H. diversifolium Broth. & Geh. —Akiyama (1986: 49); Akiyama (1988: 19); Akiyama (1989b: 159). MNP.
H. milei Mitt. subsp. **korthalsii** (Bosch & Sande Lac.) Touw —Akiyama (1986: 49); Akiyama (1988: 20); Akiyama (1989b: 159). MNP.
H. opacum M. Fleisch. —Akiyama (1986: 49, as *Hypnodendron* sp.); Akiyama (1988: 22, as *Hypnodendron* sp.); Akiyama (1989a: 199); Akiyama (1989b: 158). MNP.
H. reinwardtii (Reinw. & Hornsch.) Lindb. subsp. **caducifolium** (Herzog) Touw —Akiyama (1986: 49); Akiyama (1988: 20); Akiyama (1989b: 160). MNP.
H. subspinervium (Müll. Hal.) A. Jaeger subsp. **arborescens** (Mitt.) Touw —Akiyama (1986: 49); Akiyama (1988: 21); Akiyama (1989b: 161). MNP, NS, SS.
H. vitiense Mitt. subsp. **vitiense** —Akiyama (1986: 49); Akiyama (1988: 22); Akiyama (1989b: 161). CS, MNP.

Hypopterygiaceae

Cyathophorella (Broth.) M. Fleisch.

- C. adiantum** (Griff.) M. Fleisch. = misidentification of *Cyathophorum spinosum*, fide Akiyama (1992: 115).
C. parvifolia (Bosch & Sande Lac.) M. Fleisch. = *Cyathophorum parvifolia*, fide Kruijer (2002).
C. spinosa (Müll. Hal.) M. Fleisch. = *Cy-*

athophorum spinosum, fide Akiyama (1992: 114).

C. tenera (Bosch & Sande Lac.) M. Fleisch. = *Cyathophorum parvifolia*, fide Kruijer (2002).

C. cf. tenera (Bosch & Sande Lac.) M. Fleisch. = misidentification of *Cyathophorum parvifolia*, fide Akiyama (1992: 112).

Cyathophorum P. Beauv.

C. parvifolia Bosch & Sande Lac. —Akiyama (1986: 60, as *Cyathophorella cf. tenera*); Akiyama (1988: 42, as *Cyathophorella tenera* and *C. cf. parvifolia*); Akiyama (1992: 112, as *Cyathophorella parvifolia*); Kruijer (2002: 357).

C. spinosum (Müll. Hal.) H. Akiyama. —Akiyama (1986: 60, as *Cyathophorella spinosa* and *C. adiantum*); Akiyama (1988: 43); Akiyama (1992: 114); Kruijer (2002: 319).

Dendroclyathophorum Dixon

D. decolyi (Broth. ex M. Fleisch.) Kruijer —Akiyama (1992: 112, as *D. paradoxum*); Kruijer (2002: 290). MNP.

D. paradoxum Broth. in Cardot = *D. decolyi*, fide Kruijer 2002: 285.

Hypopterygium Brid.

H. aristatum Bosch & Sande Lac. = *H. flavolimbatum*, fide Kruijer (2002: 170).

H. cf. aristatum Bosch & Sande Lac. = misidentification of *H. flavolimbatum* and *H. vriesei*, fide Akiyama (1992: 116–117).

H. chamaedrys Bosch & Sande Lac. = *H. vriesei*, fide Kruijer (2002: 189).

H. flavolimbatum Müll. Hal. —Akiyama (1986: 60, as *H. cf. aristatum*); Akiyama (1988: 46, as *H. aristatum*); Akiyama (1992: 116, as *H. aristatum*); Kruijer (2002: 189). MNP.

Note. According to Kruijer (2002), Seram plants include both the normal and East Malesian variants.

H. humile Mitt. ex Bosch & Sande Lac. = *H. tamarisci*, fide Kruijer (2002: 200).

Note. Kruijer (2002: 187) wrote that some specimens reported as *H. humile* by Akiyama (1992) were actually *H. flavolim-*

batum.

H. tamarisci (Sw.) Brid. ex Müll. Hal. —Akiyama (1986: 60, as *H. tenellum*); Akiyama (1988: 45, as *H. humile*); Akiyama (1992: 116, as *H. humile*); Kruijer (2002: 242). MNP, NS, SS.

H. tenellum Müll. Hal. = misidentification of *H. tamarisci*, fide Akiyama (1992: 116).

H. vriesei Bosch & Sande Lac. —Akiyama (1986: 60, as *H. cf. aristatum* and *H. cf. chamaedrys*); Akiyama (1988: 46, also as *H. chamaedrys*); Akiyama (1992: 117, also as *H. chamaedrys*); Kruijer (2002: 198). CC, MNP, NS, SS, WS.

Lopidium Hook.f. & Wils.

L. javanicum Hampe. = misidentification of *L. struthiopteris*, fide Akiyama (1992: 118) and Kruijer (2002: 266).

L. struthiopteris (Brid.) M. Fleisch. —Akiyama (1986: 60, as *L. javanicum* and *L. trichocladon*); Akiyama (1988: 48, also as *L. trichocladon*); Akiyama (1992: 118, also as *L. trichocladon*); Kruijer (2002: 281). MNP, SS.

L. trichocladon (Bosch & Sande Lac.) M. Fleisch. = *L. struthiopteris*, fide Kruijer (2002: 266).

Lembophyllaceae

Camptochaete Reichdt.

C. excavata (Taylor) A. Jaeger —Akiyama (1993: 220, as *C. vaga*). MNP.

C. vaga (Müll. Hal.) Broth. ≡ *C. excavata*, fide Tangney (1997: 102)

Leptostomataceae

Leptostomum R. Brown

L. intermedium Broth. —Akiyama (1986: 48, as *L. pinaiae*); Akiyama (1988: 17); Akiyama (1989: 156). MNP.

L. pinaiae Herzog = *L. intermedium*, fide Hyvönen (1987).

Leskeaceae

Pseudoleskeopsis Broth.

P. aff. zippelii (Dozy & Molk.) Broth. —Akiyama (1986: 60).

Leucobryaceae

Leucobryum Hampe

- L. aduncum** Dozy & Molk. var. **aduncum** — Yamaguchi (1993b: 299). MNP, SS, WS.
- L. aduncum** Dozy & Molk. var. **scalare** (Müll. Hal. ex M. Fleisch.) Eddy — Akiyama (1986: 42, as *L. scarale*); Yamaguchi (1993b: 299). A, WS.
- L. arfakianum** Müll. Hal. ex Geh. — Akiyama (1986: 43, as *L. subsanctum*); Yamaguchi (1993b: 299). MNP, WS.
- L. bowringii** Mitt. — Akiyama (1986: 42); Yamaguchi (1993b: 300). MNP, WS.
- L. candidum* (P. Beauv.) Wils. var. *pentastichum* (Dozy & Molk.) Dixon = misidentification of *L. chlorophyllosum*, fide Yamaguchi (1993a).
- L. chlorophyllosum** Müll. Hal. — Akiyama (1986: 42, as *L. candidum* var. *pentastichum*); Yamaguchi (1993b: 300). CS, SS, WS.
- L. javense** (Brid. ex Schwägr.) Mitt. var. **javense** — Akiyama (1986: 42); Yamaguchi (1993b: 300). MNP, WS.
- L. javense** (Brid. ex Schwägr.) Mitt. var. **cyathifolium** (Dixon) Yamaguchi — Yamaguchi (1993b: 300). CS, MNP.
- L. javense** (Brid. ex Schwägr.) Mitt. var. **novae-guinea** (E.B. Bartram) Yamaguchi — Akiyama (1986: 42, as *L. cf. novae-guineae*); Yamaguchi (1993b: 300). MNP.
- L. cf. novae-guineae* E.B. Bartram = *L. javense* var. *novae-guinea*, fide Yamaguchi (1993a).
- L. sanctum** (Brid.) Hampe — Akiyama (1986: 42); Yamaguchi (1993b: 300). A, ES, CS, MNP, NS, SS, WS.
- L. scarale* Müll. Hal. ex M. Fleisch. = *L. aduncum* var. *scalare*, fide Yamaguchi (1993a).
- L. sericeum** Broth. ex Geh. — Yamaguchi (1993b: 301). A, WS.
- L. subsanctum* Broth. = *L. arfakianum*, fide Yamaguchi (1993a).

Schistomitrium Dozy & Molk.

- S. apiculatum** (Dozy & Molk.) Dozy & Molk. — Akiyama (1986: 43); Yamaguchi

(1993b: 301). CS, MNP.

- S. mucronifolium** (Müll. Hal.) M. Fleisch. — Akiyama (1986: 43); Yamaguchi (1993b: 301). MNP.

- S. robustum** Dozy & Molk. — Akiyama (1986: 43, as *S. cf. robustum* and *S. subrobustum*); Yamaguchi (1993b: 301). CS, MNP, WS.

- S. cf. robustum* Dozy & Molk. = misidentification of *S. robustum*, fide Yamaguchi (1993b: 301).

- S. subrobustum* Broth. = misidentification of *S. robustum*, fide Yamaguchi (1993b: 301).

Leucomiaceae

Leucomium Mitt.

- L. aneurodictyon** (Müll. Hal.) A. Jaeger — Akiyama (1986: 60); Akiyama (1989: 158). A, NS, WS.

Meteoriaceae

Aerobrydium M. Fleisch. ex Broth.

- A. crispifolium** (Broth. & Geh.) M. Fleisch. — Akiyama (1997: 9). MNP.

Aerobryopsis M. Fleisch.

- A. leptosigmata** (Geh.) M. Fleisch. — Akiyama (1986: 55); Akiyama (1997: 10). CC, MNP, SS.

- A. longissima** (Dozy & Molk.) M. Fleisch. — Akiyama (1986: 55, as *Pseudobarbella attenuata*); Akiyama (1997: 10). A, CS, MNP, NS, SS, WS.

- A. wallichii* (Brid.) M. Fleisch. = misidentification of *A. leptosigmata*, fide Akiyama (1997: 10).

Floribundaria M. Fleisch.

- F. floribunda** (Dozy & Molk.) M. Fleisch. — Akiyama (1986: 55); Akiyama (1997: 11). CS, MNP, NS, SS, WS.

- F. pseudofloribunda** M. Fleisch. — Akiyama (1986: 55); Akiyama (1997: 11). NS, CS, SS.

Meteorium Dozy & Molk.

- M. buechananii** (Brid.) Broth. — Akiyama (1997: 11). CS, MNP, NS, SS.

Note. Akiyama (1997) recognized two forms, differentiated by size, among the

Seram collections. The larger form occurs at higher elevations, the smaller one in lowland forests.

M. miquelianum (Müll. Hal.) M. Fleisch. subsp. *miquelianum* = misidentification of *M. buchananii*, fide Akiyama (1997: 11).

M. cf. subpolytrichum (Besch.) Broth. = misidentification of *M. buchananii*, fide Akiyama (1997: 11).

Papillaria A. Jaeger & Sauerb.

P. fusecescens (Hook.) A. Jaeger —Akiyama (1986: 11); Akiyama (1997:12). MNP.

Pseudobarbella Nog.

P. ancistrodes (Renauld & Cardot) Manuel —Akiyama (1997: 12). NS.

P. attenuata (Thwait. & Mitt.) Nog. = misidentification of *Aerobryopsis longissima* (Meteoriaceae), fide Akiyama (1997, 10).

Mniaceae

Plagiomnium T.J. Kop.

P. integrum (Bosch & Sande Lac.) T.J. Kop. —Akiyama (1986: 48). CS, MNP.

P. rhynchophorum (Harv.) T.J. Kop. —Akiyama (2009). CS, MNP.

P. succulentum (Mitt.) T.J. Kop. —Akiyama (1986: 48); Akiyama (2009). MNP, NS, SS.

Pohlia Hedw.

P. elongata Hedw. —Akiyama (2009). MNP.

Myuriaceae

Oedocladium Mitt.

O. rufescens (Reinw. & Hornsch.) Mitt. —Akiyama (1993: 221). MNP.

Piloecium (Müll. Hal.) Broth.

P. pseudorufescens (Hampe) Müll. Hal. in Broth. —Akiyama (1993: 221). ES, NS, WS.

Neckeraceae

Himanthocladium (Mitt.) M. Fleisch.

H. loriforme (Bosch & Sande Lac.) M. Fleisch. —Akiyama (1993: 222). MNP, NS, SS, WS.

H. plumula (Nees) M. Fleisch. —Akiyama (1993: 223). CS, MNP, NS, SS.

H. submontanum Enroth —Akiyama (1993:

223). MNP.

Homaliodendron M. Fleisch.

H. exiguum (Bosch & Sande Lac.) M. Fleisch. —Akiyama (1988: 55); Akiyama (1993: 223). ES, NS, WS.

H. flabellatum (Sm.) M. Fleisch. —Akiyama (1988: 55, as *H. scalpellifolium*); Akiyama (1993: 224). CS, MNP, SS, WS

Note. There are two distinct types in Seram, such as “scalpellifolium” type and “flabellatum” type. The former dwells in montane or mossy forests and the latter in lowland to lower montane forests.

H. microdendron (Mont.) M. Fleisch. —Akiyama (1993: 224). MNP.

Neckeropsis Reichardt

N. gracilentia (Bosch & Sande Lac.) M. Fleisch. —Akiyama (1988: 56); Akiyama (1993: 225). A, ES, MNP, NS, WS.

N. lepineana (Mont.) M. Fleisch. —Akiyama (1988: 56); Akiyama (1993: 225). CS, ES, MNP, NS, SS, WS.

Pinnatella (Müll. Hal.) M. Fleisch.

P. intralimbata M. Fleisch. —Akiyama (1988: 56); Akiyama (1993: 226). WS.

P. kuehliana (Bosch & Sande Lac.) M. Fleisch. —Akiyama (1993: 227). MNP, NS, SS, WS.

P. mariei (Besch.) Broth. —Akiyama (1993: 226). ES, NS, SS.

P. mucronata (Bosch & Sande Lac.) M. Fleisch. —Akiyama (1988: 56, as *P. cf. mucronata*); Akiyama (1993: 226). CS, MNP, NS, SS, WS.

Thamnobryum Nieuwl.

T. ellipticum (Bosch & Sande Lac.) Nog. & Z. Iwats. —Akiyama (1993: 227). MNP, SS.

T. latifolium (Bosch & Sande Lac.) Nieuwl. —Akiyama (1993: 228). MNP.

T. subserratum (Hook.) Nog. & Z. Iwats. —Akiyama (1993: 228). MNP.

T. negrosens (E.B. Bartram) Z. Iwats. & B.C. Tan —Akiyama (1993: 228); Akiyama (2009). NS, SS.

Orthotrichaceae

Desmotheca Lindb.

D. apiculata (Dozy & Molk.) Lindb. —Akiyama (1986: 50); Akiyama (2009). NS.

Groutiella Steere ex Crum & Steere

G. tomentosa (Hornsch.) Wijk & Marg. —Akiyama (2009). SS, WS.

Leptodontiopsis Broth.

L. orientalis Dixon —Akiyama (2009). MNP.

Macromitrium Brid. [identification was mostly based on Vitt *et al.* (1995) otherwise noted].

M. acuminatum (Reinw. & Hornsch.) Müll. Hal. —Akiyama (2009). MNP.

M. angustifolium Dozy & Molk. —Akiyama (2009). MNP, SS, WS.

M. archboldii E.B. Bartram —Akiyama (2009). MNP.

M. crinale Broth. & Geh. —Akiyama (2009). MNP.

M. cuspidatum Hampe —Akiyama (2009). MNP.

M. erubescens E.B. Bartram —Akiyama (2009). MNP.

M. falcatum Müll. Hal. —Akiyama (2009). SS, WS.

M. fuscescens Schwägr. —Akiyama (2009). CS, ES, MNP, NS, SS, WS.

M. incurvifolium (Hook.f. & Grev.) Schwägr. —Akiyama (2009). MNP, SS, WS.

M. longicaule Müll. Hal. —Akiyama (2009). MNP.

M. noguchianum Schultz-Motel —Akiyama (2009). MNP.

M. norrisianum Vitt. —Akiyama (2009). MNP.

M. ochraceum (Dozy & Molk.) Müll. Hal. —Akiyama (2009). MNP.

M. orthotrichum Nees ex Schwägr. —Akiyama (2009). MNP, WS.

M. parvifolium Dixon —Akiyama (2009). MNP, WS.

M. yuleanum Broth. & Geh. —Akiyama (2009). MNP.

Schlotheimia Brid.

S. emarginato-pilosa Herzog —Akiyama (2009). MNP.

S. mac-gregorii Broth. & Geh. —Akiyama (2009). MNP.

S. wallisii Müll. Hal. —Akiyama (2009). MNP.

Zygodon Hook. & Taylor

Z. intermedius Bruch & Schimp. —Akiyama (2009). MNP.

Z. novoguineensis E.B. Bartram —Akiyama (2009). MNP.

Phyllogoniaceae

Orthorrhynchium Reichardt

O. phyllogonioides (Sull.) E.B. Britton = *Cryptogonium phyllgonioides*, fide Isovita (1986).

Polytrichaceae

Dawsonia R. Brown

D. longifolia (Bruch & Schimp.) Zanten = *D. superba*, fide Zanten (1997).

D. papuana F. Müll. ex Geh. —Akiyama (1986: 39); Akiyama (1989: 152). MNP.

D. superba Grev. —Akiyama (1989: 153, as *D. longifolia*). MNP, WS.

Pogonatum P. Beauv.

P. macrophyllum Dozy & Molk. —Akiyama (1989: 154). MNP, SS, WS.

P. cf. macrophyllum Dozy & Molk. = misidentification of *P. macrophyllum*, fide Akiyama (1989).

P. neesii (Müll. Hal.) Dozy —Akiyama (1989: 156). SS, WS.

P. philippinense (Broth.) Touw —Akiyama (1989: 155). SS.

P. piliferum (Dozy & Molk.) Touw —Akiyama (1986: 39, as *Racelopus pilifer*); Akiyama (1989: 154). MNP, SS, WS.

P. subtortile (Müll. Hal.) A. Jaeger —Akiyama (1989: 155). SS.

P. cf. subtortile —Akiyama (1989: 156). MNP, WS.

Polytrichastrum G.L. Smith

P. gracile (Sw. ex Brid.) G.L. Smith —Akiyama (1986: 39, as *Polytrichum gracile*); Akiyama (1989: 153, as *P. gracile*). MNP.

Polytrichum Hedw.

P. gracile Dicks. in Menz. = *Polytrichastrum*

gracile (Sw. ex Brid.) G.L. Sm.

Racelopus Dozy & Molk.

R. pilifer Dozy & Molk. = *Pogonatum pilifer*,
fide Touw (1986: 29).

Plagiotheciaceae

Plagiothecium Bruch & Schimp.

P. cf. euryphyllum (Cardot & Thér.) Z. Iwats.
—Akiyama (1988: 52). MNP.

Pottiaceae

Anoetangium Schwägr.

A. aestivum (Hedw.) Mitt. —Akiyama (1996:
158). MNP.

Barbula Hedw.

B. consanguinea (Thwait. & Mitt.) A. Jaeger
—Akiyama (1996: 159). CS, ES.

B. javanica Dozy & Molk. —Akiyama (1996:
161). ES, NS, SS, WS.

B. novoguinese Broth. —Akiyama (1996:
163). NS.

B. pachyloma Broth. —Akiyama (1996: 163);
Akiyama (2009). A, MNP, NS, SS, WS.

B. robbinsii E.B. Bartram —Akiyama (1996:
165). MNP, SS.

***B. seramensis** H. Akiyama —Akiyama
(1996: 161). NS, SS.

Type: Central Seram, en route from Nihe-
hata to Hatumete, 180 m alt., *Akiyama*
C-16743 (holotype in HYO, isotypes in
BM, BO, FH, H, L, MO, NY).

Gymnostomum Nees & Hornsch.

G. aurantiacum (Mitt.) Paris —Akiyama
(1996: 165). CS, MNP, SS, WS.

Hyophila Brid.

H. involuta (Hook.) A. Jaeger —Akiyama
(1996: 165). ES, NS, SS, MNP, WS.

Leptodontium (Müll. Hal.) Lindb.

L. flexifolium (Dicks. ex With.) Hampe —
Akiyama (1996: 166). MNP.

Pseudosymblepharis Broth.

P. angustata (Mitt.) Hilp. —Akiyama (1996:
166). MNP.

Splachnobryum Müll. Hal.

S. limbatum Norris & Zander —Akiyama
(2009). SS.

S. obtusum (Brid.) Müll. Hal. —Akiyama

(2009). SS.

Trichostomum Bruch

T. brachydontium Bruch ex F.A. Müll. —
Akiyama (1996: 166). MNP.

Weissia Hedw.

W. edentula Mitt. —Akiyama (1996: 166).
CS.

W. aff. exerta (Broth.) Chen —Akiyama
(1996: 167). MNP.

Pterobryaceae

Calypothecium Mitt.

C. integrifolium (E.B. Bartram) Nog. —Aki-
yama (1993: 250). SS.

C. recurvulum (Broth.) Broth. in Thér. —
Akiyama (1996: 54, as *C. tumidum*); Aki-
yama (1990: 136). CS, NS, WS.

C. tumidum (Dicks. ex Hook.) M. Fleisch. =
C. recurvulum, fide Hyvönen (1989).

C. aff. urvilleanum (Müll. Hal.) Broth. —
Akiyama (1993: 250). MNP.

Cryptogonium (Müll. Hal.) Hampe

C. phyllgonioides (Sull.) Isov. —Akiyama
(1986: 55, as *Orthorrhynchium phyllogoni-*
oides); Akiyama (1990: 137). CS, NS, SS,
WW.

Euptychium Schimp.

Euptychium sp. = misidentification of *Ga-*
rovaglia compressa var. *compressa* (Ga-
rovagliaceae), fide Akiyama (1990: 140).

Neolindbergia

N. vitiensis (E.B. Bartram) Enroth —Akiya-
ma (1986: 53, as *Forsstoremia* sp.); Akiya-
ma *et al.* (1991: 82). MNP, SS.

N. veloirae H. Akiyama —Akiyama *et al.*
(1991: 84). CS.

Pterobryopsis

P. crassicaulis (Müll. Hal.) M. Fleisch. —
Akiyama (1986: 54, as *P. aff. gedehensis*);
Akiyama (1990: 137). MNP.

P. aff. gedehensis M. Fleisch. = misidentifica-
tion of *P. crassicaulis*, see Akiyama (1990,
137).

Symphysodon Dozy & Molk.

S. neckeroides Dozy & Molk. —Akiyama
1990: 137 (also as *S. subneckeroides*). CS,
MNP, SS, WS.

S. subneckeroides Broth. = misidentification of *Symphyodon neckeroides*, see Akiyama 1990: 137.

Symphysodontella M.Fleisch.

S. attenuata (Mont.) M. Fleisch. —Akiyama (1986: 54, as *S. cf. attenuata*); Akiyama (1990: 138). MNP.

S. cf. subulata Broth. —Akiyama (1986: 54, as *Symphysodontella* sp.); Touw (1992: 350).

S. parviflora E.B. Bartram —Akiyama (1986: 54); Akiyama (1990: 138). MNP.

S. cylindracea (Mont.) M. Fleisch. —Akiyama (1986: 54); Akiyama (1990: 138). SS, MNP, WS.

S. splendens (Reinw. & Hornsch.) Touw & Magill —Akiyama (1986: 54, as *S. cf. obtusa*); Akiyama (1990: 138). NS.

S. cf. obtusa Tixier = misidentification of *S. splendens*, see Touw (1992: 350).

S. sp. = misidentification of *S. cf. subulata*, see Touw (1992: 350).

Ptychomniaceae

Glyphothecium Hampe

G. sciuiroides (Hook.) Hampe —Akiyama (1986: 50, as *G. aff. sciuiroides*); Akiyama (1989: 157). MNP.

G. aff. sciuiroides (Hook.) Hampe = misidentification of *G. sciuiroides*, fide Akiyama (1989: 157).

Racopilaceae

Powellia Mitt.

**P. macromitrioides* H. Akiyama —Akiyama (1988: 24, as *Powellia* sp.); Akiyama (1990: 130). NS (MNP).

Type: Indonesia [Moluccas]. Seram: Rumah Sokat Batu near Sawai, Kecamatan Seram Utara, 230 m, Akiyama C-9930 (holotype in KYO, isotypes in BO, NY).

P. parvula (E.B. Bartram) Norris & T.J. Kop. —Akiyama (2009). MNP.

Powellia sp. = misidentification of *P. macromitrioides*, see Akiyama (1990: 130).

Racopilum P. Beauv.

R. cuspidigerum (Schwägr.) Aongstr. —Akiyama (1986: 53); Akiyama (1988: 27); Akiyama (1990: 133). MNP, NS, SS, WS.

Note. Akiyama (1988) recognized two types in this pantropical species; one corresponding to typical *R. cuspidigerum*, while the other has upright capsules, extended pilifera at leaf apices, and much smaller laminal cells.

R. spectabile Reinw. & Hornsch. —Akiyama (1986: 53); Akiyama (1988: 26); Akiyama (1990: 134). CS, MNP, NS, SS, WS.

R. aff. spectabile —Akiyama (1988: 26). MNP.

R. verrucosum Herzog —Akiyama (1986: 53); Akiyama (1988: 28); Akiyama (1990: 135). MNP.

Note. This rare species was originally described based on Seram specimens (Herzog 1916: 238). Akiyama & van Zanten (1999) reported another species with strongly verrucose setae, *Racopilum niutensis* Akiyama & Zanten, from West Kalimantan.

Regmatodontaceae

Regmatodon Brid.

R. cf. polycarpus (Griff.) Mitt. —Akiyama (2009). MNP.

Rhizogoniaceae

Hymenodon Hook.f. & Wils.

H. angustifolius Sande Lac. —Akiyama (1990: 130). MNP.

H. pilifer Hook.f. & Wils. subsp. *sericeus* (Dozy & Molk.) Kartt. & Bäck —Akiyama (1986: 48, as *H. sericeus*); Akiyama (1990: 130). MNP.

H. cf. sriceus (Dozy & Molk.) Müll. Hal. = misidentification of *H. pilifer* subsp. *sericeus*, fide Akiyama (1990: 130) and Koponen *et al.* (1986).

Hymenodontopsis Herzog

H. stresemanii Herzog —Akiyama (1986: 48); Akiyama (1990: 129). MNP.

Pyrrhobryum Mitt.

P. latifolium (Bosch. & Sande Lac.) Mitt. —

Akiyama (1990: 131). CS.

P. spiniforme (Hedw.) Mitt. —Akiyama (1986: 49); Akiyama (1990: 131). CS, MNP, SS, WS.

Rhizogonium Brid.

R. graeffeanum (Müll. Hal.) A. Jaeger —Akiyama (1990: 130). MNP.

Sematophyllaceae

Acanthorrhynchium M. Fleisch.

A. papillatum (Harv.) M. Fleisch. —Akiyama (1993: 231). NS, ES.

Acroporium Mitt.

A. diminutum (Brid.) M. Fleisch. —Akiyama (1993: 232, also as *A. subulatum*). CS, MNP, WS.

A. hamulatum (M. Fleisch.) M. Fleisch. —Akiyama (1993: 232). MNP.

Note. This species was collected from Seram Island under the name of *A. warburgii* (Broth.) M. Fleisch. in L & NY.

A. hermaphroditum (Müll. Hal.) M. Fleisch. —Akiyama (1993: 232). MNP, WS.

A. longicaule (Sande Lac.) M. Fleisch. —Akiyama (1993: 233). MNP, WS.

A. secundum (Reinw. & Hornsch.) M. Fleisch. —Akiyama (1993: 233). MNP.

Note. Specimens referred to *A. stramineum* (Reinw. & Hornsch.) M. Fleisch. were collected from Ambon (*Robinson 2269 & 2279*, NY!), but they are actually *A. secundum*.

A. aff. secundum —Akiyama (1993: 233). WS.

A. sigmatodontium (Müll. Hal.) M. Fleisch. —Akiyama (1993: 233). CS, MNP, WS.

Acroporium sp. —Akiyama (1993: 233). MNP.

Aptychella (Broth.) Herzog

A. scalaris (Müll. Hal.) M. Fleisch. —Akiyama (1993: 233). MNP.

Aptychella sp. = misidentification of *Dimorphocladon borneense* (Symphyodontaceae), fide Akiyama (1997: 29).

Brotherella Loeske ex M. Fleisch.

B. falcata (Dozy & Molk.) M. Fleisch. —Akiyama (1993: 235). MNP.

Clastobryum Dozy & Molk.

C. cf. epiphyllum (Renauld & Cardot) B.C. Tan —Akiyama (1993: 236). MNP, SS.

Glossadelphus M. Fleisch.

G. zollingeri (Müll. Hal.) M. Fleisch. —Akiyama (1993: 236). NS.

G. aff. zollingeri —Akiyama (1993: 236). WS.

Isocladiella Dixon

I. surcularis (Dixon) B.C. Tan & Mohamed —Akiyama (1993: 236, as *I. phyllogonioides*). MNP, WS.

I. phyllogonioides Dixon = *I. surcularis*, fide Tan & Mohamed (1990).

Macrohymenium Müll. Hal.

M. rufum Müll. Hal. —Akiyama (1993: 237). MNP, SS.

Mastopoma Cardot

M. armittii (Broth. & Geh.) Broth. —Akiyama (1993: 238). MNP.

M. brauniana (Bosch & Sande Lac.) H. Akiyama —Akiyama (1993: 238); Akiyama (2006: 9). MNP.

M. aff. robinsonii (Broth.) E.B. Bartram —Akiyama (1993: 238). MNP.

Meiothecium Mitt.

M. aff. bogoriense M. Fleisch. —Akiyama (2009). WS.

M. hamatum (Müll. Hal.) Broth. —Akiyama (1993: 239); Akiyama (2009). MNP, WS.

M. jagorii (Müll. Hal.) Broth. —Akiyama (1993: 239). WS.

M. microcarpum (Hook.) Mitt. —Akiyama (1993: 240); Akiyama (2009). A, SS, WS.

Papillidiopsis (Broth.) W.R. Buck & B.C. Tan

P. luxurians (Dozy & Molk.) W.R. Buck & B.C. Tan —Akiyama (1993: 240). NS.

Pseudohypnella (Broth.) M. Fleisch.

P. verrucosa (Bosch & Sande Lac.) M. Fleisch. —Akiyama (1993: 241); Akiyama (2009). MNP.

Rhaphidostichum M. Fleisch.

R. bunodiocarpum (Müll. Hal.) M. Fleisch. —Akiyama (1993: 241). ES, MNP, NS.

Rhaphidostichum sp. —Akiyama (1993: 242). A, WS.

Sematophyllum Mitt.

S. microcladiellum M. Fleisch. —Akiyama (1993: 242). A, ES, NS, WS.

S. saproxylophilum (Müll. Hal.) M. Fleisch. —Akiyama (1993: 242). SS, WS.

Taxithelium Spruce

***T. damanhurianum** P.S. Câmara —Câmara (2009: 590); Akiyama (2009). MNP, NS.

Type: Northern Seram, Manusela national Park, Rumah Sokat Batu in the vicinity of Sawai, 100 m alt., *Akiyama C-9923* (holotype in HYO; isotypes in L, NY).

Note. Câmara (2009) designated *Akiyama C-9329* as the type of *T. damanhurianum*, but it must be a typographical error. Among the three specimens (*C-9329*, *C-9409* and *C-9923*) that he identified as *T. damanhurianum*, only *C-9923* has complete sporophytes. In addition, judging from information given for the type locality, as well as annotation on the specimens, *Akiyama C-9923* must be the type of *T. damanhurianum*.

T. isoclada (Bosch & Sande Lac.) Renauld & Cardot —Akiyama (2009). SS.

T. kerianum (Broth.) Broth. —Akiyama (1993: 243). MNP.

T. levieri (Broth. & Geh.) Broth. —Akiyama (1993: 243). MNP, NS.

T. lindbergii (Sande Lac.) Renauld & Cardot —Akiyama (1993: 244). NS.

T. nepalense (Schwägr.) Broth. = misidentification of *T. planum*, fide Akiyama (1993: 244).

T. planum (Brid.) Mitt. —Akiyama (1993: 244). ES, CS, MNP, NS, SS, WS.

Note. We confirmed the presence of *T. planum* in Ambon (*Robinson 2272 & 2311* in L and NY; *Zippelius s.n.*, in L).

Trichosteleum Mitt.

T. hamatum (Dozy & Molk.) A. Jaeger —Akiyama (1993: 245). CS, ES, MNP, NS, SS, WS.

Note. We confirmed the presence of *T. hamatum* in Ambon (*Robinson 2283 & 2319* in FH, L and NY).

T. aff. hamatum —Akiyama (1993: 246). NS, SS, WS.

T. mannosum (Müll. Hal.) A. Jaeger —Akiyama (1993: 246). MNP, SS.

Trismegistia (Müll. Hal.) Broth.

T. calderensis (Sull.) Broth. —Akiyama (1986: 64, as *T. cf. rigida* and *T. cf. rigida f. pendula*); Akiyama (1993: 247, as *T. rigida*). MNP, WS.

T. lancifolia (Harv.) Broth. —Akiyama (1993: 247). CS, MNP, NP, SS, WS.

Note. Both var. *lancifolia* and var. *korthalsii* (Dozy & Molk.) M. Fleisch. are present in Seram Island. We also confirmed the presence of *T. lancifolia* in Ambon (*Robinson 2280*, FH).

T. rigida (Mitt.) Broth. = misidentification of *T. calderensis*, fide Akiyama (1993: 247).

T. cf. rigida (Mitt.) Broth. = misidentification of *T. calderensis*, fide Akiyama (1993: 247).

T. cf. rigida (Mitt.) Broth. f. *pendula* M. Fleisch. = misidentification of *T. calderensis*, fide Akiyama (1993: 248).

T. panduriformis (Wright) Broth. —Akiyama (1986: 64, as *T. cf. panduriformis*); Akiyama (1993: 248). MNP.

T. cf. panduriformis (Wright) Broth. = misidentification of *T. panduriformis*, fide Akiyama (1993: 248).

Warbugeilla Müll. Hal.

W. aureo-nitida (E.B. Bartram) H. Akiyama —Akiyama (1993: 249). MNP.

W. breviseta (Broth.) Broth. —Akiyama (1993: 249). MNP.

W. subleptorrhynchoides (M. Fleisch.) M. Fleisch. —Akiyama (1993: 249). MNP.

Sorapillaceae

Sorapilla Spruce & Mitt.

S. papuana Broth. & Geh. in Broth. —Akiyama (1989: 157); Akiyama (2009). MNP.

Sphagnaceae

Sphagnum L.

S. cuspidatum Ehrh. in Hoffm. var. **cuspidatum** —Akiyama (1989: 151). SW.

Spiridentaceae

Spiridens Nees

S. longifolius Lindb. —Akiyama (1988: 24, as *S. reinwardtii*); Akiyama (1989: 158, as *S. reinwardtii*). MNP.

S. reinwardtii Nees = misidentification of *S. longifolius*.

Splachnaceae

Tayloria Hook.

T. indica Mitt. —Akiyama (1986: 47, as *T. aff. longiseta*); Akiyama (1993: 220); Akiyama (2009). MNP.

T. aff. longiseta E.B. Bartram = misidentification of *T. indica*, fide Akiyama (1993: 220).

Symphyodontaceae

Chaetomitrium Dozy & Molk.

C. acanthocarpum Bosch & Sande Lac. —Akiyama (1986: 57); Akiyama (1997: 19). NS, SS.

C. elegans Geh. —Akiyama (1986: 57, as *C. cf. tahitense*); Akiyama (1997: 20). ES, NS, WS.

C. cf. longisetulum E.B. Bartram = misidentification of *C. subplicatum*, fide Akiyama (1997:26) and Akiyama (1999: 259).

C. leptopoma (Schwägr.) Bosch & Sande Lac. = misidentification of *C. spinosum*, fide Akiyama (1997: 25).

C. orthorrhynchum (Dozy & Molk.) Bosch & Sande Lac. —Akiyama (1986: 57, as *C. cf. orthorrhynchum*); Akiyama (1997: 20). NS, SS, WS.

C. cf. orthorrhynchum (Dozy & Molk.) Bosch & Sande Lac. = misidentification of *C. orthorrhynchum*, fide Akiyama (1997: 20).

C. papillifolium Bosch & Sande Lac. —Akiyama (1986: 57); Akiyama (1997: 22). NS.

C. perlaeve Dixon —Akiyama (1997: 22). MNP.

C. cf. perlaeve Dixon = misidentification of *C. perlaeve*, fide Akiyama (1997: 22).

***C. seramensis** H. Akiyama —Akiyama (1997: 26, as *C. aff. spinosum*); Akiyama (1999: 258). SS.

Type: Central Seram, en route from Wae

Heka Heka to Wae Pasola Hatu, ca. 500 m alt., *Akiyama C-16011* (holotype in HYO, isotypes in BO, L).

C. spinosum E.B. Bartram —Akiyama (1986: 57, as *C. leptopoma*); Akiyama (1986: 57, as *C. cf. spinosum*); Akiyama (1997: 25, as *C. cf. spinosum*); Akiyama (1999: 259). MNP.

C. cf. spinosum E.B. Bartram = misidentification of *C. spinosum*, fide (Akiyama 1999: 259).

C. aff. spinosum E.B. Bartram = misidentification of *C. seramensis*, fide Akiyama (1999: 258).

C. subplicatum E.B. Bartram —Akiyama (1986: 57, as *C. cf. longisetulum*); Akiyama (1997: 26, as *C. aff. subplicatum*); Akiyama (1999: 259). NS.

C. aff. subplicatum E.B. Bartram = misidentification of *C. subplicatum*, fide Akiyama (1999: 259).

C. cf. tahitense (Sull.) Mitt. = misidentification of *C. elegans*, fide Akiyama (1997: 20).

C. torquescens Bosch & Sande Lac. —Akiyama (1997: 27). NS, SS.

C. virescens Bosch & Sande Lac. —Akiyama (1986: 57); Akiyama (1997: 20). CS, MNP, NS, SS.

Dimorphocladon Dixon

D. borneense Dixon —Akiyama (1993: 235, as *Aptychella* sp.); Akiyama (1997: 29); Akiyama (2009). CS, NS.

Thuidiaceae

Aequatoriella Touw

A. bifaria (Bosch & Sande Lac.) Touw —Akiyama (1986: 61, as *Pelekium bifarium*); Watanabe & Akiyama (1991: 65, as *Pelekium bifarium*); Touw (1993: 197, as *Thuidium hattorii*).

Claopodium (Lesq. & James) Renauld & Cardot

C. cf. assurgens (Sull. & Lesq.) Cardot = misidentification of *C. prionophyllum*, fide Watanabe & Akiyama (1991: 65).

C. prionophyllum (Müll. Hal.) Broth. —Akiyama (1986: 61, as *C. cf. assurgens*); Watanabe & Akiyama (1991: 65). MNP.

Indothuidium Touw

- I. kiaense** (R.S. Williams) Touw —Akiyama (1986: 62, as *Thuidium* aff. *kiaense*); Watanabe & Akiyama (1991: 62, as *Thuidium kiaense*).

Pelekium Mitt.

- P. bifarium* (Bosch & Sande Lac.) M. Fleisch. = *Aequatoriella bifaria*, fide Touw (2001b: 202).
- P. bonianum** (Besch.) Touw —Akiyama (1986: 62, as *T. meyenianum*); Watanabe & Akiyama (1991: 62, as *Thuidium bonianum*). WS.
- P. gratum** (P. Beauv.) Touw —Watanabe & Akiyama (1991: 63, as *Thuidium kuripanum*). ES, MNP, NS, SS, WS.
- P. investe** (Mitt.) Touw —Akiyama (1986: 62, as *Thuidium tamariscellum*); Watanabe & Akiyama (1991: 62, as *T. investe* and *T. tamariscellum*). MNP, SS.
- P. velatum** Mitt. —Akiyama (1986: 61); Watanabe & Akiyama (1991: 625). CS, NS, SS, WS.
- P. synoicum** (Touw) Touw —Touw (2001a: 117).
- P. watanabei** (Touw) Touw —Watanabe & Akiyama (1991: 62, as *Thuidium contortulum*); Touw (1993: 200, as *T. watanabei*); Touw (2001a: 87). MNP.

Thuidium Bruch & Schimp.

- T. bonianum* Besch. = *Pelekium bonianum*, fide Touw (2001b: 203).
- T. contortulum* (Mitt.) A. Jaeger = misidentification of *Pelekium watanabei* (= *T. watanabei*; Thuidiaceae), fide Touw (1993: 202).
- T. cymbifolium** (Dozy & Molk.) Dozy & Molk. —Akiyama (1986: 61); Watanabe & Akiyama (1991: 62). MNP, NN, SS.
- T. glaucinoides* Broth. = *T. pristocalyx* var. *samoanum*, fide Touw (2001).
- T. glaucinum* (Mitt.) Bosch & Sande Lac. = *T. pristocalyx* var. *samoanum*, fide Touw (2001).
- T. hattorii* Touw = *Aequatoriella bifaria*, fide Touw (2001a: 63).
- T. investe* (Mitt.) A. Jaeger = *Pelekium investe*, fide Touw (2001b: 203).

T. kuripanum (Dozy & Molk.) R. Watanabe = *Pelekium gratum*, fide Touw (2001a: 107).

T. kiaense R.S. Williams = *Indothuidium kiaense*, fide Touw (2001a: 122 and his note 1 on p. 124).

T. aff. kiaense R.S. Williams = misidentification of *Indothuidium kiaense* (Thuidiaceae), fide Watanabe & Akiyama (1991: 63).

T. kuripanum (Dozy & Molk.) R. Watanabe = *Pelekium gratum*, fide Touw (2001a: 107)

T. meyenianum (Hampe) Dozy & Molk. = misidentification of *P. bonianum* (Thuidiaceae), fide Watanabe & Akiyama (1991: 62) and Touw (2001a: 113).

T. plumulosum (Dozy & Molk.) Dozy & Molk. —Akiyama (1986: 62); Watanabe & Akiyama (1991: 64).

T. pristocalyx (Müll. Hal.) A. Jaeger var. **pristocalyx** —Touw (2001a, fig. 8, without citation of specimens). unknown.

T. pristocalyx var. **samoanum** (Mitt.) Touw —Akiyama (1986: 62, as *T. glaucinoides* and *T. glaucinum*); Watanabe & Akiyama (1991: 64, as *T. samoanum*); Touw (2001b, fig. 10, without citation of specimens examined). CC, MNP, NS, WS

T. samoanum Mitt. = *T. pristocalyx* var. *samoanum*, fide Touw (2001a: 41. See also his notes on p. 30).

T. tamariscellum (Müll. Hal.) Bosch & Sande Lac. = misidentification of *Pelekium investe* (Thuidiaceae), fide Touw (2001: 104).

T. watanabei Touw = *Pelekium watanabei*, fide Touw (2001b: 205).

Trachylomataceae

Trachyloma Brid.

T. indicum Mitt. —Akiyama (1896: 5); Akiyama 1990: 139. CC, MNP, SS, WS.

Trachypodaceae

Trachypus Reinw. & Hornsch.

T. bicolor Reinw. & Hornsch. —Akiyama (1986: 53, as *T. bicolor* and *Trachypus* sp.)

T. aff. appendiculatus (Renauld & Cardot) Broth. —Akiyama (1986: 53) reported on the basis of two specimens, but later Aki-

yama (1997) did not listed this species from the Islands.

Trachypus sp. = misidentification of *T. bicolor*, fide Akiyama (1986: 53)

HEPATICAEE

(Aneuraceae are not included.)

Adelanthaceae

Wettsteinia Schiffn.

W. inversa (Sande Lac.) Schiffn. —Kitagawa (1986: 67). MNP.

Alisoniaceae

Sandeothallus R.M. Schust.

S. radiculosus (Schiffn.) R.M. Schust. —Kitagawa (1986: 71). MNP, SS.

Balantiopsaceae

Balantiopsis Mitt.

B. ciliaris S. Hatt. subsp. **novoguineensis** S. Hatt. —Kitagawa (1986: 70). MNP.

Calypogeiaceae

Calypogeia Raddi

C. fusca (Lehm.) Steph. —Kitagawa (1986: 67). CS.

C. goebelii (Schiffn.) Steph. —Kitagawa (1986: 67). MNP.

Cephaloziaceae

Odontoschisma (Dumort.) Dumort.

Odontoschisma sp. —Kitagawa (1986: 67). MNP.

Frullaniaceae

Frullania Raddi

***F. akiyamae** S. Hatt. —Hattori (1986: 240). MNP.

Type: Central Seram, Tehoru, from Hatumete to Holae Pass, 1430 m alt., *Akiyama C-10675* (holotype in NICH, isotype in KYO).

F. apiculata (Reinw. *et al.*) Dumort. —Hattori (1986: 241). CS, MNP, SS,

F. armatifolia Verd. —Hattori (1986: 242). MNP.

F. attenuata Steph. —Hattori (1986: 242).

MNP.

***F. cordistipula** (Reinw. *et al.*) Dumort. var. **dentistipula** S. Hatt. —Hattori (1986: 242). MNP.

Type: Central Seram, Manusela National Park, Kanikeh, along a trail between Wae Angsela and Wae Huhu, 1480 m alt., *Akiyama C-8761* (holotype in NICH, isotype in KYO).

F. ericoides (Nees) Mont. —Hattori (1986: 242). CS.

F. gaudichaudii (Nees & Mont.) Nees & Mont. —Hattori (1986: 242). MNP.

F. gracilis (Reinw. *et al.*) Dumort. var. **gracilis** —Hattori (1986: 242). ES, MNP.

***F. gracilis** var. **vittata** S. Hatt. —Hattori (1986: 243). MNP.

Type: North Seram, Sawai, Mt. Musisi, 1300 m alt., *Akiyama C-9788* (holotype in NICH, isotype in KYO).

F. hasskarliana Lindenb. var. **hasskarliana** —Hattori (1986: 243). MNP.

***F. hasskarliana** var. **gracilis** S. Hatt. —Hattori (1986: 243). NS.

Type: Central Seram, trail between Hatumete and Hoale Pass, 1400m alt., *Akiyama C-10731* (holotype in NICH, isotype in KYO).

***F. hasskarliana** var. **parvidentata** S. Hatt. —Hattori (1986: 245). MNP.

Type: Central Seram, Manusela National Park, trail between Wae Angsela and Wae Huhu, 1000 m alt., *Akiyama C-8800* (holotype in NICH, isotype in KYO).

F. heteromorpha Schiffn. —Hattori (1986: 245). NS.

F. junghuhniana Gottsche var. **junghuhniana** —Hattori (1986: 245). MNP.

F. junghuhniana var. **tenella** (Sande Lac.)

- Grolle & S. Hatt. —Hattori (1986: 245). MNP.
- F. multilacera** Steph. —Hattori (1986: 245). NS, WS.
- F. nobilis** Steph. var. **cochleata** (Steph.) S. Hatt. —Hattori (1986: 246). MNP.
- F. nodulosa** (Reinw. *et al.*) Nees —Hattori (1986: 246). NS, WS.
- F. orientalis** Sande Lac. —Hattori (1986: 246). CS.
- F. pauciramea** Steph. —Hattori (1986: 246). MNP.
- ***F. piptophylla** S. Hatt. var. **minor** S. Hatt. —Hattori (1986: 247). MNP.
Type: Central Seram, Manusela National Park, trail between Wae Huhu and G. O'Wae Puku, 2520 m alt., *Akiyama C-8937* (holotype in NICH, isotype in KYO).
- F. pulogensis** Steph. —Hattori (1986: 247). MNP.
- F. ramuligera** (Nees.) Mont. —Hattori (1986: 247). MNP.
- F. reflexistipula** Sande Lac. —Hattori (1986: 247). MNP.
- F. repandistipula** Sande Lac. —Hattori (1986: 247). MNP.
- ***F. serrata** Gottsche var. **ceramensis** S. Hatt. —Hattori (1986: 247). MNP.
Type: Central Seram, Manusela National Park, Mt. Musisi near Sawai, *Akiyama C-9788* (holotype in NICH, isotype in KYO).
- F. serrata** var. **hamatispina** (S. Hatt.) S. Hatt. —Hattori (1986: 248). MNP.
- F. serrata** fo. **crispulodentata** Verd. —Hattori (1986: 247). MNP.
- F. subcaduca** S. Hatt. —Hattori (1986: 248). MNP.
- F. sublignosa** Steph. —Hattori (1986: 248). ES, WS.
- F. subnigricaulis** S. Hatt. var. **subtruncata** S. Hatt. —Hattori (1986: 248). MNP.
- ***F. suocellata** S. Hatt. —Hattori (1986: 248). —Hattori (1986: 248). MNP.
Type: Central Seram, trail between Hatumete and Hoale Pass, 1030 m alt., *Aki-*

yama C-10631 (holotype in NICH, isotype in KYO).

- F. ternatensis** Gottsche var. **non-appendiculata** S. Hatt. —Hattori (1986: 248). MNP.

Geocalycaceae

Conoscyphus Mitt.

- C. trapezioides** (Sande Lac.) Schiffn. —Kitagawa (1986: 68). MNP.

Heteroscyphus Schiffn.

- H. argutus** (Reinw. *et al.*) Schiffn. —Kitagawa (1986: 69). ES, NS, SS.
- H. aselliformis** (Reinw. *et al.*) Schiffn. —Kitagawa (1986: 69). MNP.
- H. coalitus** (Hook.) Schiffn. —Kitagawa (1986: 69). MNP.
- H. splendens** (Lehm. & Lindenb.) Grolle —Kitagawa (1986: 69). NS, MNP.
- H. zollingeri** (Gottsche) Schiffn. —Kitagawa (1986: 69). SS.

Lophocolea (Dumort.) Dumort.

- L. aff. ciliolata** (Nees) Gottsche —Kitagawa (1986: 69). NS, MNP.
- L. aff. erustiana** (Steph.) N. Kitag. —Kitagawa (1986: 69). MNP.
- L. giulianettii** Steph. —Kitagawa (1986: 69). SS.
- L. kurzii** Sande Lac. —Kitagawa (1986: 69). NS.

Saccogynidium Grolle

- S. muricellum** (De Not.) Grolle —Kitagawa (1986: 69). MNP.

Herbertaceae

Herbartus Gray

- H. armitanus** (Steph.) H.A. Mill. —Kitagawa (1986: 65); Juslén (2006: 415). MNP.
- H. javanicus** (Steph.) H.A. Mill. —Kitagawa (1986: 65). MNP.
- H. longifissus** Steph. —Kitagawa (1986: 65). MNP.
- H. longispinus** Jack & Steph. —Kitagawa (1986: 65). MNP.

Jungermanniaceae

Anastrophyllum (Spruce) Steph.

- A. bidens** (Nees) Steph. —Kitagawa (1986:

68). MNP.

A. revolutum Steph. —Kitagawa (1986: 68). MNP.

Anomacaulis (R.M. Schust.) Grolle

A. flaccidus (Steph.) Grolle —Kitagawa (1986: 68). MNP.

Chandonanthus Mitt.

C. hirtellus (Web.) Mitt. —Kitagawa (1986: 68). MNP.

Denotarisia Grolle

D. linguifolia (De Not.) Grolle —Kitagawa (1986: 68). MNP.

Jamesoniella (Spruce) Carrington

J. contracta (Reinw. *et al.*) N. Kitag. —Kitagawa (1986: 68). CS.

J. flexicaulis (Nees) Schiffn. —Kitagawa (1986: 68). MNP.

Jungermannia L.

J. ariadne Tayl. —Kitagawa (1986: 68). WS.

J. tetragona Lindenb. —Kitagawa (1986: 68). CS, SS.

J. truncata Nees —Kitagawa (1986: 68). A, E.

Syzygiella Spruce

S. ovalifolia Inoue —Kitagawa (1986: 68). MNP.

S. securifolia (Nees) Inoue —Kitagawa (1986: 68). MNP.

Lejeuneaceae

Acrolejeunea (Spruce) Schiffn.

A. pycnoclada (Tayl.) Schiffn. —Mizutani (1986: 299). NS, SS.

A. securifolia (Endl.) Watts —Mizutani (1986: 299). MNP.

Caudalejeunea Steph.

C. circinata Steph. —Mizutani (1986: 299). CS, MNP, SS.

C. reniloba (Gottsche) Steph. —Mizutani (1986: 299). ES, MNP, NS, SS.

Ceratolejeunea (Spruce) Schiffn.

C. belongeriana (Gottsche) Steph. —Mizutani (1986: 299). CS, NS.

C. moniliata Herzog —Mizutani (1986: 299). CS, MNP, SS.

Cheilolejeunea (Spruce) Schiffn.

C. ceylanica (Gottsche) R.M. Schust. & Ka-

chr. —Mizutani (1986: 300). MNP.

C. imbricata (Nees) S. Hatt. —Mizutani (1986: 300). CS, MNP, SS.

C. intertexta (Lindenb.) Steph. —Mizutani (1986: 300). NS.

C. meyeniana (Gottsche *et al.*) R.M. Schust. —Mizutani (1986: 300). MNP.

C. occlusa (Herzog) Kodama & N. Kitag. —Mizutani (1986: 300). MNP, SS.

C. vittata (Steph.) Steph. —Mizutani (1986: 300). CS, ES, MNP, NS, SS.

Cololejeunea (Spruce) Schiffn.

C. crenulata (Herzog) Benedix. —Mizutani (1986: 300). CS, MNP, SS.

C. dozyana (Sande Lac.) Schiffn. —Mizutani (1986: 300); Zhu & So (1998: 125). MNP.

C. equalbi Tixier —Mizutani (1986: 300). MNP, NS, SS, WS.

C. falcatooides Benedix. —Mizutani (1986: 300). CS, SS.

C. floccosa (Lehm. & Lindenb.) Schiffn. —Mizutani (1986: 300). ES, CS, NS, WS.

C. goebelii (Schiffn.) Schiffn. —Mizutani (1986: 301). CS, ES, MNP, NS, SS, SW.

C. hasskarliana (Lehm. & Lindenb.) Schiffn. —Mizutani (1986: 301). MNP.

C. javanica (Steph.) Mizut. —Mizutani (1986: 301). CS, MNP.

C. lanciloba Steph. —Mizutani (1986: 301). CS, MNP, NS, SS, WS.

C. nymanii (Steph.) Benedix —Mizutani (1986: 301). MNP, SS, WS.

C. papillosa (K.I. Goebel) Mizut. —Mizutani (1986: 301). CS, MNP.

C. peculiaris (Herzog) Benedix —Mizutani (1986: 301). CS, MNP, SS.

C. plagiophylla Benedix —Mizutani (1986: 301). CS, NS.

C. planissima (Mitt.) Abyw. —Mizutani (1986: 301). CS, MNP.

C. raduliloba Steph. —Mizutani (1986: 301). SS.

C. stephanii Benedix —Mizutani (1986: 301). MNP.

C. triapiculata (Herzog) Tixier —Mizutani (1986: 301). MNP.

C. yulensis (Steph.) Benedix —Mizutani

(1986: 302). CS, MNP.

Colura (Dumort.) Dumort.

C. acroloba (Mont.) Jovet-Ast —Mizutani (1986: 302). MNP, WS.

C. ari (Steph.) Steph. —Mizutani (1986: 302). CS, MNP, NS.

C. conica (Sande Lac.) Grolle —Mizutani (1986: 302). CS, MNP.

C. imperfecta Steph. —Mizutani (1986: 302). MNP, SS.

C. ornata K.I. Goebel —Mizutani (1986: 302). SS.

C. superba (Mont.) Steph. —Mizutani (1986: 302). SS.

Diplasiolejeunea (Spruce) Schiffn.

D. cavifolia Steph. —Mizutani (1986: 302). CS.

D. patelligera Herzog —Mizutani (1986: 302). MNP.

Drepanolejeunea (Spruce) Schiffn.

D. dactylophora (Gottsche *et al.*) Schiffn. —Mizutani (1986: 302). CS, MNP.

D. grandis Herzog —Mizutani (1986: 302). MNP.

D. intermedia Zwickel —Mizutani (1986: 302). CS, MNP.

D. levicornua Steph. var. **levicornua** —Mizutani (1986: 303). NS, MNP, SS.

D. levicornua var. **longicornua** Herzog —Mizutani (1986: 303). MNP, NS.

D. lyrata Grolle —Mizutani (1986: 303). MNP.

D. micholitzii Steph. —Mizutani (1986: 303). CS, MNP.

D. spinosocornuta Steph. —Mizutani (1986: 303). MNP.

D. tenera K.I. Goebel —Mizutani (1986: 303). MNP, SS.

D. ternatensis (Gottsche) Steph. —Mizutani (1986: 303). CS, MNP, SS.

Hattoriolejeunea Mizut.

***H. akiyamae** Mizut. —Mizutani (1986: 303). MNP.

Type: Central Seram, Tehoru, Hatumete —Hoale Pass, 1160–1760 m alt., *Akiyama C-10650* (holotype in KYO, isotype NICH).

Lejeunea Lib.

L. albescens (Steph.) Mizut. —Mizutani (1986: 305). CS, MNP.

L. catanduana (Steph.) Mill. *et al.* —Mizutani (1986: 305). NS.

L. cucullata (Reinw. *et al.*) Nees —Mizutani (1986: 305). CS, MNP, NS.

L. exilis (Reinw. *et al.*) Grolle —Mizutani (1986: 305); Zhu & Grolle (2003: 102). CS, MNP, NS.

L. fissistipula (Steph.) Steph. —Mizutani (1986: 306). CS, MNP, NS.

L. lumbricoides (Nees) Nees —Mizutani (1986: 306). MNP.

L. microloba Tayl. —Mizutani (1986: 306). MNP.

L. sordida (Nees) Nees —Mizutani (1986: 306). ES, SS.

Lepidolejeunea R.M. Schust.

L. bidentula (Steph.) Steph. —Mizutani (1986: 306). NS.

L. dentistipula Steph. —Mizutani (1986: 306). NS.

L. elliptica (Lehm & Lindenb.) Schiffn. —Mizutani (1986: 306). ES, NS, WS.

L. epiphylla (Mitt.) Steph. —Mizutani (1986: 306). CS, NS, MNP, WS.

L. foliicola Steph. —Mizutani (1986: 306). MNP.

L. maculata (Mitt.) Steph. —Mizutani (1986: 306). MNP, NS, MNP, SS.

Lopholejeunea (Spruce) Schiffn.

L. eulopha (Tayl.) Schiffn. —Mizutani (1986: 306). SS.

L. horticola Schiffn. —Mizutani (1986: 306). ES, SS.

L. javanica (Nees) Schiffn. —Mizutani (1986: 306). ES, SS.

L. nigricans (Lindenb.) Schiffn. —Mizutani (1986: 307). NS.

L. subfusca (Nees) Schiffn. —Mizutani (1986: 307). CS, ES, MNP, NS, WS.

Mastigolejeunea (Spruce) Schiffn.

M. humilis (Gottsche) Schiffn. —Mizutani (1986: 307). ES, NS, WS.

M. ligulata (Lehm. & Lindenb.) Schiffn. —Mizutani (1986: 307). ES, NS.

M. truncata Mizut. —Mizutani (1986: 307). NS.

M. undulata Gradst. & Grolle —Mizutani (1986: 307). MNP, NS, SS, WS.

M. virens (Aongstr.) Steph. —Mizutani (1986: 307). MPN, NS, WS.

Metzgeriopsis K.I. Goebel

M. pusilla K.I. Goebel —Mizutani (1986: 307). MNP.

Otolejeunea Grolle & Tixier

O. semperiana (Steph.) Grolle —Mizutani (1986: 307). NS.

Ptychanthus Nees

P. striatus (Lehm. & Lindenb.) Nees —Mizutani (1986: 307). WS.

Rhaphidolejeunea Herzog

R. longicruris (Steph.) Herzog —Mizutani (1986: 307). CS, MNP.

Schiffneriolejeunea Verd.

S. tumida (Nees) Gradst. —Mizutani (1986: 308). WS.

Spruceanthus Verd.

S. marianus (Gottsche) Mizut. —Mizutani (1986: 308). NS.

S. polymorphus (Sande Lac.) Verd. —Mizutani (1986: 308). CS, MNP, NS, SS.

Thysananthus Lindenb.

T. appendiculatus Steph. —Mizutani (1986: 308). MNP, SS.

T. convolutus Lindenb. —Mizutani (1986: 308). CS, NS.

T. fruticosus (Lindenb. & Gottsche) Schiffn. —Mizutani (1986: 308). NS, SS.

T. spatulistipus (Reinw. *et al.*) Lindenb. —Mizutani (1986: 308). CS, NS, SS.

Tuyamaella S. Hatt.

T. serratistipa S. Hatt. —Mizutani (1986: 308). MNP.

Lepicoleaceae

Lepicolea Dumort.

L. rara (Steph.) Grolle —Kitagawa (1986: 65). MNP.

L. yakushimensis (S. Hatt.) S. Hatt. —Kitagawa (1986: 65). MNP.

Mastigophora Nees

M. diclados (Brid.) Nees —Kitagawa (1986:

65). NS, CS, MNP, SS.

Lepidoziaceae

Acromastigum A. Evans

A. divaricatum (Gottsche *et al.*) Evans —Kitagawa (1986: 65). MNP.

Arachniopsis Spruce

A. major Herzog —Kitagawa (1986: 65). NS.

Bazzania Gray

B. australis (Mont.) Trev. —Kitagawa (1986: 66). NS, SS.

B. cadens N. Kitag. —Kitagawa (1986: 66). NS.

B. cinnamata (De Not.) Trev. —Kitagawa (1986: 66). MNP.

B. commutata (Lindb. & Gottsche) Schiffn. —Kitagawa (1986: 66). MNP.

B. conophylla (Sande Lac.) Steph. —Kitagawa (1986: 66). NS.

B. distans (Nees) Trev. —Kitagawa (1986: 66). MNP.

B. erosa (Reinw. *et al.*) Trev. —Kitagawa (1986: 66). CS, MNP, NS.

B. insignis (De Not.) Trev. —Kitagawa (1986: 66). MNP.

B. kernii Steph. —Kitagawa (1986: 66). MNP.

B. kokawana N. Kitag. & Kodama —Kitagawa (1986: 66). MNP.

B. longicaulis (Sande Lac.) Schiffn. —Kitagawa (1986: 66). MNP.

B. loricata (Reinw. *et al.*) Trev. —Kitagawa (1986: 66). MNP.

B. parvixta Steph. —Kitagawa (1986: 66). CS, MNP.

B. pectinata (Lindenb. & Gottsche) Schiffn. —Kitagawa (1986: 66). MNP.

B. spiralis (Reinw. *et al.*) Meijer —Kitagawa (1986: 66). MNP.

B. tridens (Reinw. *et al.*) Trev. —Kitagawa (1986: 66). NS, MNP.

B. vittata (Gottsche) Trev. —Kitagawa (1986: 66). MNP, SS.

Kurzia G. Martens

K. geniculata Mizut. —Kitagawa (1986: 66). CS.

K. gonyotricha (Sande Lac.) Grolle —Kitagawa (1986: 66). SS.

Lepidozia (Dumort.) Dumort.

L. borneensis Steph. —Kitagawa (1986: 66);
Doei (1987b: 518). MNP.

L. brotheri Steph. —Doei (1987b: 520). MNP.

L. cladorhiza (Reinw. *et al.*) Nees —Kitagawa
(1986: 67); Doei (1987b: 520). MNP.

L. ferdinandii-muelleri Steph. —Kitagawa
(1986: 67); Doei (1987b: 521). MNP.

L. hasskarliana Steph. —Kitagawa (1986:
67); Doei (1987b: 521). MNP.

Note. Doei (1987: 528) noted that *Lepido-
zia ceramensis* Herzog (Herzog 1926: 340)
might be conspecific to *L. hasskarliana*
Steph.

***L. integrifolia** Doei —Doei (1987a: 421).
MNP, WS.

Type: West Seram, Tihulale, en route from
the upper elevation to the top of Gunung
(Mt.) Totaniwel, ca. 1170 m alt., *Akiya-
ma C-15811* (holotype in KYO, isotype
in BO).

L. papulosa Steph. —Kitagawa (1986: 67);
Doei (1987b: 522). CS, MNP, NS, SS, WS.

L. subintegra Lindenb. = misidentification of
L. subtrichodes, fide Doei (1987: 525).

L. subtrichodes Steph. —Kitagawa (1986: 67
as *L. subintegra*); Doei (1987b: 523). MNP.

L. trichodes (Reinw. *et al.*) Nees —Kitagawa
(1986: 67); Doei (1987b: 526). CS, MNP,
SS, WS.

L. wallichiana Gottsche —Doei (1987b: 527).
MNP, WS.

Psiloclada Mitt.

P. clandestina Mitt. —Kitagawa (1986: 67).
CS.

Telaranea Spruce ex Schiffn.

T. kogiana (Setph.) Grolle —Kitagawa (1986:
67). NS.

T. neesii (Lindenb.) Fulford —Kitagawa
(1986: 67). MNP.

T. trisetosa (Steph.) Grolle —Kitagawa
(1986: 67). MNP.

Zoopsis (Hook.f. & Taylor) Gottsche *et al.*

Z. liukiensis Horik. —Kitagawa (1986: 67).
MNP.

Z. setigera K.I. Goebel —Kitagawa (1986:
67). CS.

Metzgeriaceae

Metzgeria Raddi

M. foliicola Schiffn. —Kitagawa (1986: 71).
MNP.

M. leptoneura Spruce —Kitagawa (1986:
71). CS, NS, SS.

Pallaviciniaceae

Podomitrium Mitt.

P. malaccense (Spruce) D. Campb. —Kitaga-
wa (1986: 71). NS.

Plagiochilaceae

Plagiochila (Dumort.) Dumort.

***P. akiyamae** Inoue —Inoue (1986: 73). MNP.

Type: Central Seram, Tehoru, along a trail
between Hatumete and Holae Pass,
200–640 m alt., *Akiyama C-10532* (ho-
lotype in TNS, isotype in KYO).

P. arbuscula (Brid. ex Lehm. & Lindenb.)
Lindenb. —Inoue (1986: 76). MNP.

P. bantamensis (Reinw. *et al.*) Dumort. —In-
oue (1986: 76). CS.

***P. ceramica** Inoue —Inoue (1986: 76). MNP.
Type: Manusela National Park, around the
summit of G. O'Wae Puku, 2800–2900
m alt., *Akiyama C-8992* (holotype in
TNS, isotype in KYO).

P. decidua Inoue & Grolle —Inoue (1986:
78). MNP.

P. dendroides (Nees) Lindenb. —Inoue
(1986: 78). CS, MNP, NS.

P. frondescens (Nees) Lindenb. —Inoue
(1986: 78). MNP.

P. gedeana Schiffn. —Inoue (1986: 78). MNP.

P. massalongoana Schiffn. —Inoue (1986:
79). NS, SS, WS.

P. novae-guineae Sande Lac. —Inoue (1986:
79). NS.

P. pseudaberrans Inoue & Grolle —Inoue
(1986: 81). MNP.

P. renitens (Nees) Lindenb. —Inoue (1986:
81). MNP.

P. salacensis Gottsche —Inoue (1986: 81). ES,
MNP.

P. sandei Dozy —Inoue (1986: 81). CS.

P. sciophila Nees in Lindenb. —Inoue (1986:

83). MNP.

P. seemannii Mitt. —Inoue (1986: 83). CS, MNP, NS, SS.

P. teysmanii Sande Lac. —Inoue (1986: 84). MNP, NS.

Plagiochilion S. Hatt.

P. aff. fimbriatum (Mitt.) Inoue —Kitagawa (1986: 69). CS.

P. oppositus (Reinw. *et al.*) S. Hatt. —Kitagawa (1986: 69). MNP.

Pleuroziaceae

Pleurozia Dumort.

P. gigantea (Web.) Lindb. —Kitagawa (1986: 70). MNP.

Pseudolepicoleaceae

Temnoma Mitt. & Hook.f.

T. setigerum (Lindenb.) R.M. Schust. —Kitagawa (1986: 65). MNP.

Radulaceae

Radula Dumort.

R. anceps Sande Lac. —Kitagawa (1986: 70). MNP, NS, SS.

R. campanigera Mont. —Kitagawa (1986: 70). MNP.

R. densifolia Castel —Kitagawa (1986: 70). CS.

R. javanica Gottsche —Kitagawa (1986: 70). NS.

R. nymanii Steph. —Kitagawa (1986: 70). MNP.

R. sumatrana Steph. —Kitagawa (1986: 70).

CS.

R. tjibodensis K.I. Goebel —Kitagawa (1986: 70). MNP, NS, SS

R. vanzantienii Yamada —Kitagawa (1986: 70). NS.

R. vrieseana Sande Lac. —Kitagawa (1986: 70). ES, NS, WS.

Schgistochilaceae

Schistochila (Nees) Schiffn.

S. aligera (Nees *et al.*) Jack & Steph. —Kitagawa (1986: 69). CS.

S. formosana Horik. —Kitagawa (1986: 69). MNP.

S. philippinensis (Mont.) Jack & Steph. —Kitagawa (1986: 69). CS, NS, MNP.

S. recurvata Buch —Kitagawa (1986: 69). MNP.

S. sciurea (Nees) Schiffn. —Kitagawa (1986: 69). MNP.

S. aff. schultzei Steph. —Kitagawa (1986: 69). MNP.

Treubiaceae

Treubia K.I. Goebel

T. insignis K.I. Goebel —Kitagawa (1986: 70). MNP.

Trichocoleaceae

Trichocolea Dumort.

T. breviseta Steph. —Kitagawa (1986: 65). MNP.

T. pluma (Reinw. *et al.*) Mont. —Kitagawa (1986: 65). MNP, NS, SS.

ANTHOCEROTAE

Anthocerotaceae

Dendroceros Mees

D. acutilobus Steph. —Hasegawa (1986: 9). NS, MNP.

D. cavernosus Haseg. —Hasegawa (1986: 10). CS, NS.

D. foliicola Haseg. —Hasegawa (1986: 10). CS.

D. javanicus (Nees) Nees in Gottsche —

Hasegawa (1986: 10). MNP.

***D. seramensis** Haseg. —Hasegawa (1986: 10). NS, SS.

Type: Central Seram, Tehoru, between Wolu and Wae Waya, 0–360 m alt., *Akiyama C-10269* (KYO).

Folioceros D.C. Bhardwaj

F. appendiculatus (Steph.) Haseg. —Hasegawa (1986: 13). NS.

Megaceros Campb.

M. flagellaris (Mitt.) Steph. —Hasegawa (1986: 13). CS, MNP, SS.

Phaeoceros Prosk.

P. laevis (L.) Prosk. subsp. **caroliniamus**

(Michaux) Prosk. —Hasegawa (1986: 13). ES, NS, WS.

P. polyandrus (Steph.) Haseg. —Hasegawa (1986: 13). A.

I thank to Dr. Masahiro Kato (National Museum of Nature and Science, Tokyo), Dr. Dedy Darnaedi (LIPI) and other members of the expeditions to the islands. Identification of specimens are due to the assistance of the following authorities; Dr. P. E. A. S. Câmara (*Taxithelium*), Dr. H. Doei (Lepidoziaceae), Dr. J. Hasegawa (Anthocerotae), Dr. S. Hattori (Frullaniaceae), Dr. H. Inoue (Plagiogochilaceae), Dr. Z. Iwatsuki and Dr. T. Suzuki (Fissidentaceae), Dr. N. Kitagawa (general hepatics), Dr. H. Kruijer (Hypopterygiaceae), Dr. M. Mizutani (Lejeuneaceae), Dr. A. Touw (Thuidiaceae), Dr. J. Vana (Jungermanniaceae), Dr. D. Vitt (Calomniaceae) and Dr. T. Yamaguchi (Leucobryaceae and Leucophanaceae).

References

- Akiyama, H. 1986. Bryophyte Flora. In: Kato, M. (ed.) Taxonomic studies of the plants of Seram Island, pp. 36–74. Botanical Garden, University of Tokyo, Tokyo.
- Akiyama, H. 1988. Bryophyte. In: Kato, M. (ed.), Taxonomic Studies of the Plants of Seram Island, pp. 13–59. Botanical Garden, University of Tokyo, Tokyo.
- Akiyama, H. 1989a. A reevaluation of *Hypnodendron opacum* (Hypnodendraceae, Musci). Bryologist 92: 198–201.
- Akiyama, H. 1989b. Taxonomic studies of mosses of Seram and Ambon (Moluccas, East Indonesia) collected by Indonesian-Japanese Botanical Expeditions I. Acta Phytotax. Geobot. 40: 147–162.
- Akiyama, H. 1990. Taxonomic studies of mosses of Seram and Ambon (Moluccas, East Indonesia) collected by Indonesian-Japanese Botanical Expeditions II. J. Fac. Sci., Univ. Tokyo, Bot. 14: 385–413.
- Akiyama, H. 1990. Taxonomic studies of mosses of Seram and Ambon (Moluccas, East Malesia) collected by Indonesian-Japanese Botanical Expedition III. Acta Phytotax. Geobot. 41: 129–142.
- Akiyama, H. 1992. Taxonomic studies of mosses of Seram and Ambon (Moluccas, East Malesia) collected by Indonesia-Japanese Botanical Expeditions, V. Hypopterygiaceae. Acta Phytotax. Geobot. 43: 111–120.
- Akiyama, H. 1993. Taxonomic studies of mosses of Seram and Ambon (Moluccas, East Malesia) collected by Indonesian-Japanese Botanical Expeditions. VII, (Michaux) Prosk. —Hasegawa (1986: 13). ES, NS, WS.
- P. polyandrus** (Steph.) Haseg. —Hasegawa (1986: 13). A.
- Lembophyllaceae, Splachnaceae, Myuriaceae, Nekeraceae, Brachytheciaceae, Amblystegiaceae and Sematophyllaceae, with additions to previous reports on Dicranaceae and Pterobryaceae. J. Fac. Sci., Univ. Tokyo, Bot. 15: 219–254.
- Akiyama, H. 1996. Taxonomic studies of mosses of Seram and Ambon (Moluccas, East Malesia) collected by Indonesian-Japanese Botanical Expedition, IX. Pottiaceae. Trop. Bryol. 12: 157–168.
- Akiyama, H. 1997a. Taxonomic studies of mosses of Seram and Ambon (Moluccas, East Malesia) collected by Indonesian-Japanese Botanical Expeditions VIII. Meteoriaceae, Hookeriaceae, and Trachypodaceae. Nat. Human Activ. 2: 9–31.
- Akiyama, H. 1997b. [Abstract] Bryophyte flora of Seram Island, East Malesia. IAB Symposium on 2000's Bryology, Abstracts: 103.
- Akiyama, H. 1999. Additional notes on the genus *Chaetomitrium* Dozy & Molk. (Hookeriaceae, Musci) reported from Seram Island (Moluccas, East Malesia). Bryol. Res. 7: 257–260.
- Akiyama, H. 2006. Notes on the taxa previously classified in the genus *Trismegistia* (Sematophyllaceae). J. Hattori Bot. Lab. 100: 1–14.
- Akiyama, H. 2009. Taxonomic studies of mosses of Seram and Ambon (Moluccas, East Malesia) collected by Indonesian-Japanese Botanical Expeditions X. Bartramiaceae, Brachytheciaceae, Bryaceae, Ditrichaceae, Erpodiaceae, Orthotrichaceae, and additions and corrections to the previous reports. Nat. Human Activ. 20 [in press].
- Akiyama, H., T. Koponen & D. H. Norris. 1991. Bryophyte flora of the Huon Peninsula, Papua New Guinea. XLV. *Neolindbergia* (Prionodontaceae, Musci). Acta Bot. Fennica 143: 71–89.
- Akiyama, H. & W. D. Reese. 1993. Taxonomic studies of mosses of Seram and Ambon (Moluccas, East Malesia) collected by Indonesian-Japanese botanical expeditions. VI. Calymperaceae. J. Fac. Sci., Univ. Tokyo, Bot. 15: 199–218.
- Akiyama, H. & M. Suleiman. 2001. Taxonomical notes on the genus *Chaetomitrium* (Hookeriaceae, Musci) of Borneo. Hikobia 13: 491–509.
- Akiyama, H. & B. O. van Zanten. 1999. *Racopilum niutensis*, a new species from Kalimantan Barat (Indonesian Borneo) and notes on Bornean species of the Ra-

- copilaceae (Bryopsida). *Bryobrothera* 5: 17–21.
- Bruggeman-Nannenga, M. A. 1997. Notes on *Fissidens* VI. New synonyms, new combinations and validation of some names. *J. Hattori Bot. Lab.* 81: 155–173.
- Câmara, P. E. A. S. (2009). A new species of *Taxithelium* (Sematophyllaceae) from Seram, Indonesia. *Bryologist* 112: 589–592.
- Chien, G. (ed). 1994. *Flora Bryophytorum Sinicorum* 1. Beijing.
- Crosby, M. R., U. Schultze-Motel & W. Schultze-Motel. 1983. Katalog der Laubmoose von Madagaskar und den umliegenden Inseln. *Willdenowia* 13: 187–255.
- Doei, H. 1987a. A new species of *Lepidozia* (Hepaticae) from Seram Island, Indonesia. *J. Hattori Bot. Lab.* 63: 421–424.
- Doei, H. 1987b. The genus *Lepidozia* (Hepaticae) from Seram Island, Indonesia. In: *Plant Ecology and Taxonomy to the Memory of Dr. Satoshi Nakanishi*, pp. 517–528. The Kobe Geobotanical Society, Kobe.
- Dozy, F. & J. H. Molkenboer. 1844. *Muscorum Frondosorum Novae Species ex Archipelago Indico et Japonica*. Leiden.
- Dozy, F. & J. H. Molkenboer. 1845–1854. *Musci Frondosi Inediti Archipelagi Indici*. Leiden.
- Dozy, F. & J. H. Molkenboer. 1854–1861. *Bryologia Javanica*, 1. Leiden.
- Dozy, F. & J. H. Molkenboer. 1861–1870. *Bryologia Javanica*, 2. Leiden.
- Fleischer, M. 1900–1922. *Die Musci der Flora von Buitenzorg*, 4 vols. Leiden.
- Hasegawa, J. 1986. A collection of the Anthocerotae from Seram and Ambon. *Acta Phytotax. Geobot.* 37: 9–15.
- Hattori, S. 1986. *Frullania* collection made by Mr. Akiyama on Seram Island. *J. Hattori Bot. Lab.* 60: 239–253.
- Herzog, T. 1916. Neue laubmoose aus Ostasien und Südamerika. *Hedwigia* 57: 233–250.
- Herzog, T. 1919. Die Laubmoose der II. Freiburger Mokukkenexpedition. *Hedwigia* 61: 286–302.
- Herzog, T. 1926. Bryophyten der weiteren Indomalaya. *Hedwigia* 66: 337–358.
- Hofmann, H. 1997. A monograph of the genus *Palamodinium* (Brachytheciaceae, Musci). *Lindbergia* 22: 3–20.
- Hyvönen, J. 1987. A synopsis of the genus *Leptostomum* (Leptostomataceae, Musci). *Ann. Bot. Fennici* 24: 63–72.
- Hyvönen, J. 1989. Bryophyte flora of the Huon Peninsula, Papua New Guinea. XXVI. Pterobryaceae (Musci). *Acta Bot. Fennica* 137: 1–40.
- Isoviita, P. 1986. *Cryptogonium* replaces *Pursellia* (Musci, Pterobryaceae). *J. Hattori Bot. Lab.* 60: 451–453.
- Inoue, H. 1986. Contributions to the knowledge of the Plagiochilaceae of southeastern Asia XIX. The Genus *Plagiochila* from Isl. Seram. *Bull. Nat. Sci. Mus., Ser. B, Bot.* 12, 73–84.
- Iwatsuki, Z. & T. Suzuki. 1988. *Fissidens* collections made by Mr. H. Akiyama on Seram Island in 1985 and 1986. *Hikobia* 10: 215–220.
- Juslén, A. 2006. Revision of Asian *Herbertus* (Herbertaceae, Marchantiophyta). *Ann. Bot. Fenn.* 43: 409–436.
- Kitagawa, N. 1986. Hepaticae. In: Kato, M. (ed.), *Taxonomic Studies of the Plants of Seram Island*, pp. 65–71. Botanical Garden, University of Tokyo, Tokyo.
- Klazenga, N. 1999. A revision of the Malesian species of *Dicranoloma* (Dicranaceae, Musci). *J. Hattori Bot. Lab.* 87: 1–130.
- Koponen, T., A. Touw & D. H. Norris. 1986. Bryophyte flora of the Huon Peninsula, Papua New Guinea. XIV. Rhizogoniaceae (Musci). *Acta Bot. Fennica* 133: 1–24.
- Kruijer, H. 2002. Hypopterygiaceae of the world. *Blumea, Supplement* 13: 1–388.
- Magombo, Z. 2003. Taxonomic revision of the moss family Diphysciaceae M. Fleisch. (Musci). *J. Hattori Bot. Lab.* 94: 1–86.
- Mizutani, M. 1986. Lejeuneaceae from Seram Island, Indonesia. *J. Hattori Bot. Lab.* 61: 299–308.
- Mohamed, H. & W. D. Reese. 1985. *Syrrhopodon* (Musci: Calymperaceae) in Malaysia and adjacent regions. *Bryologist* 88: 223–254.
- Norris, D. H. & T. Koponen. 1998. Bryophyte flora of the Huon Peninsula, Papua New Guinea. XXIV Andreaeaceae, Archidiaceae, Seligeriaceae, Ditrichaceae (Musci). *Ann. Bot. Fennici* 25: 165–177.
- Pippo, S. 1984. Bryophyte flora of the Huon Peninsula, Papua New Guinea. III. Haplomitriaceae, Lepicoleaceae, Herbertaceae, Pseudlepicoleaceae, Trichocoleaceae, Schistochilaceae, Balantiopsidaceae, Pleuroziaceae and Porellaceae (Hepaticae). *Ann. Bot. Fennici* 21: 21–48.
- Reese, W. R., H. Akiyama & C. C. Townsend 1991. *Syrrhopodon pulcher*, a new species from Seram and Sulawesi. *Bryologist* 94: 419–420.
- Salazar Allen, N. 1993. A revision of the pantropical genus *Leucophanes* Brid. *Bryophyt. Biblioth.* 46: 1–281.
- Schiffner, V. 1893. Über exotische Hepaticae, hauptsächlich aus Java, Amboina und Brasilien, nebst einigen morphologischen und krisischen Bemerkungen über *Marchantia*. *Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.* 60: 219–316.
- Schiffner, V. 1910. Über die Gattungen *Chiloscyphus* und *Heteroscyphus* n. gen. *Oesterr. Bot. Zeitschr.* 60: 169–173.
- Stephani, F. 1898–1924. *Species Hepaticarum*, I–VI. Georg & Cie, Genève et Bale.
- Tan, B. C. & T. Koponen. 1983. *Dicranoloma* (Musci, Dicranaceae) in Southeast Asia, with special reference to the Philippine taxa. *Ann. Bot. Fennici* 20: 317–334.
- Tangney, R. S. 1997. A taxonomic revision of the genus *Camptochaete* Reichdt., Lembophyllaceae (Musci). *J.*

- Hattori Bot. Lab. 81: 53–121.
- Touw, A. 1986. A revision of *Pogonatum* sect. *Racelopus*, sect. nov., including *Racelopus* Dozy & Molk., *Pseudoracelopus* Broth., and *Racelopodopsis* Thér. J. Hattori Bot. Lab. 60: 1–60.
- Touw, A. 1992. A survey of the mosses of the Lesser Sunda Islands (Nusa Tenggara), Indonesia. J. Hattori Bot. Lab. 71: 289–366.
- Touw, A. 1993. Notes on tropical Asian Thuidiaceae, with two new Malesian species of *Thuidium* s.l. J. Hattori Bot. Lab. 74: 193–204.
- Touw, A. 2001a. A review of the Thuidiaceae (Musci) and a realignment of taxa traditionally accommodated in *Thuidium* sensu amplo (*Thuidium* Schimp., *Thuidiopsis* (Broth.) M. Fleisch., and *Pelekium* Mitt.) including *Aequatoriella* gen. nov., and *Indothuidium* gen. nov. J. Hattori Bot. Lab. 90: 167–209.
- Touw, A. 2001b. A taxonomic revision of the Thuidiaceae (Musci) of tropical Asia, the western Pacific, and Hawaii. J. Hattori Bot. Lab. 91: 1–136.
- Vitt, D. H. 1995. The genus *Calomnion* (Bryopsida): taxonomy, phylogeny, and biogeography. Bryologist 98: 338–358.
- Watanabe, R. & H. Akiyama. 1991. Taxonomic studies of mosses of Seram and Ambon (Moluccas, East Malaysia) collected by Indonesian-Japanese Botanical Expedition IV. Thuidiaceae. Acta Phytotax. Geobot. 21: 61–66.
- Yamaguchi, T. 1993a. A revision of the genus *Leucobryum* (Musci) in Asia. J. Hattori Bot. Lab. 73: 1–123.
- Yamaguchi, T. 1993b. Leucobryaceae and Leucophanoideae (Calymperaceae) on Seram and Ambon Islands. Hikobia 11: 299–303.
- Zanten, B. O. van. 1997. Proposal to conserve the name *Dawsonia superba* (Musci, Dawsoniaceae). Taxon 46: 547–549.
- Zanten, B. O. van. 2006. A synoptic review of the Racopilaceae (Bryophyta, Musci). 1. Asian, Pacific and Australasian species of the genus *Racopilum*. J. Hattori Bot. Lab. 100: 527–552.
- Zhu, R.-L. & R. Grolle. 2003. Taxonomy and distribution of *Lejeunea exilis* (Lejeuneaceae, Hepaticae). Ann. Bot. Fenn. 40: 101–106.
- Zhu, R.-L. & M.-L. So. 1998. Two epiphyllous liverworts, *Cololejeunea dozyana* (Sande Lac.) Schiffn. and *Cololejeunea sigmoidea* Jovet-Ast & Tixier (Hepaticae, Lejeuneaceae), new to Taiwan. Bot. Bull. Acad. Sin. 39: 125–130.

Received July 22, 2009; accepted August 31, 2009